

A 2-COLORING OF $[1, N]$ CAN HAVE $(1/22)N^2 + O(N)$ MONOCHROMATIC SCHUR TRIPLES, BUT NOT LESS!

AARON ROBERTSON AND DORON ZEILBERGER¹

Department of Mathematics, Temple University, Philadelphia, PA 19122, USA

aaron@math.temple.edu, zeilberg@math.temple.edu

Submitted: March 3, 1998; Accepted: March 25, 1998; Revised: May 3, 1998

Abstract: WE PROVE THAT THE MINIMUM NUMBER (ASYMPTOTICALLY) OF MONOCHROMATIC SCHUR TRIPLES THAT A 2-COLORING OF $[1, n]$ CAN HAVE IS $\frac{n^2}{22} + O(n)$. THIS REVISED VERSION FILLS IN A MINOR AND SUBTLE GAP DISCOVERED BY M. PRIMAK. (THE REVISION ALSO CORRECTS (AT NO EXTRA COST) A DISCREPANCY BETWEEN THE SOLUTION IN THE PAPER AND THE SOLUTION OBTAINED BY MAPLE. IN THE PAPER $H_{1/2}$ SHOULD BE H_0 AND H_1 SHOULD BE $H_{1/2}$ FOR THE SOLUTIONS TO AGREE.)

This article is accompanied by the Maple package RON, available from either author's website.

Tianjin, June 29, 1996: In a fascinating invited talk at the SOCA 96 combinatorics conference organized by Bill Chen, Ron Graham proposed (see also [GRR], p. 390):

Problem (\$100): Find (asymptotically) the least number of monochromatic Schur triples $\{i, j, i + j\}$ that may occur in a 2-coloring of the integers $1, 2, \dots, n$.

By renaming the two colors 0 and 1, the above is equivalent to the following

Discrete Calculus Problem: Find the minimal value of

$$F(x_1, \dots, x_n) := \sum_{\substack{1 \leq i < j \leq n \\ i+j \leq n}} [x_i x_j x_{i+j} + (1 - x_i)(1 - x_j)(1 - x_{i+j})],$$

over the n -dimensional (discrete) unit cube $\{(x_1, \dots, x_n) | x_i = 0, 1\}$. We will determine *all* local minima (with respect to the Hamming metric), then determine the global minimum.

Partial Derivatives: For any function $f(x_1, \dots, x_n)$ on $\{0, 1\}^n$ define the discrete *partial derivatives* $\partial_r f$ by $\partial_r f(x_1, \dots, x_r, \dots, x_n) := f(x_1, \dots, x_r, \dots, x_n) - f(x_1, \dots, 1 - x_r, \dots, x_n)$.

If (z_1, \dots, z_n) is a local minimum of F , then we have the n inequalities:

$$\partial_r F(z_1, \dots, z_n) \leq 0 \quad , \quad 1 \leq r \leq n.$$

A purely routine calculation (applicable Maple routines: `diff1`, `dif`) shows that (below $\chi(S)$ is 1(0) if S is true(false))

$$\partial_r F(x_1, \dots, x_n) =$$

¹ Supported by NSF.

Mathematics Classification Numbers: Primary: 05D10, 05A16; Secondary: 04A20

$$(2x_r-1) \left\{ \sum_{i=1}^n x_i + \sum_{i=1}^{n-r} x_i - (n - \lfloor \frac{r}{2} \rfloor) - \chi(r > \frac{n}{2}) - (2x_r-1) + x_r \chi(r > \frac{n}{2}) + 1 - (x_{\frac{r}{2}} + x_{2r}) \chi(r \leq \frac{n}{2}) \right\}.$$

Since we are only interested in the *asymptotic* behavior, we can modify F by any amount that is $O(n)$. In particular, we can replace $F(x_1, \dots, x_n)$ by

$$G(x_1, \dots, x_n) = F(x_1, \dots, x_n) + \sum_{i=1}^{n/2} x_i (x_{2i} - 1) - \frac{1}{2} \sum_{i=1}^n x_i.$$

Noting that $(2x_r - 1)^2 \equiv 1$ and $(2x_r - 1)x_r \equiv x_r$ on $\{0, 1\}^n$, we see that for $1 \leq r \leq n$,

$$\partial_r G(x_1, \dots, x_n) = (2x_r - 1) \left\{ \sum_{i=1}^n x_i + \sum_{i=1}^{n-r} x_i - (n - \lfloor \frac{r}{2} \rfloor) - \frac{1}{2} \chi(r \leq n/2) \right\} - \frac{1}{2} \chi(r \leq n/2) - 1/2.$$

Let $k = \sum_{i=1}^n x_i$. Since at a local minimum (z_1, \dots, z_n) we have $\partial_r G(z_1, \dots, z_n) \leq 0$, it follows that any local minimum (z_1, \dots, z_n) satisfies the

Ping-Pong Recurrence: Choose $a, b \in \{0, 1\}$ arbitrarily each time \hat{H} or \tilde{H} is used, where \hat{H} and \tilde{H} are the following functions:

$$\hat{H}(y) := \begin{cases} 0, & \text{if } y > 1/2; \\ 1, & \text{if } y < 0; \\ a, & \text{if } 0 \leq y \leq 1/2. \end{cases}$$

$$\tilde{H}(y) := \begin{cases} 0, & \text{if } y > 1; \\ 1, & \text{if } y < -1; \\ b, & \text{if } -1 \leq y \leq 1. \end{cases}$$

Then we must have, for $r = n, n-1, \dots, n - \lfloor n/2 \rfloor + 1$,

$$z_r = \hat{H} \left(k - n + \lfloor \frac{r}{2} \rfloor + \sum_{j=1}^{n-r} z_j \right), \quad (\text{Right Volley})$$

$$z_{n-r+1} = \tilde{H} \left(2k - n - 1/2 + \lfloor \frac{n-r+1}{2} \rfloor - \sum_{j=r}^n z_j \right), \quad (\text{Left Volley})$$

and if n is odd then $z_{(n+1)/2} = \hat{H}(k - n + \lfloor \frac{n+1}{4} \rfloor + \sum_{j=1}^{(n-1)/2} z_j)$.

These equations determine a solution (depending upon the choices of the a 's and b 's made along the way), z (if it exists), in the order $z_n, z_1, z_{n-1}, z_2, \dots$. When we solve the Ping-Pong recurrence we forget the fact that $\sum_{i=1}^n z_i = k$. Most of the time a solution will not satisfy this last condition, but when it does, we have a genuine local minimum. Note that *any* local minimum must show up in this way.

Solutions of the Ping-Pong Recurrence: By playing around with the Maple routine `ptor2` in our Maple package `RON` (available from either author's website), we were able to find the following

solutions, for n sufficiently large, to the Ping-Pong recurrence. As usual, for any word (or letter) W , W^m means ‘ W repeated m times’.

Let $w = 2k - n$, $k \neq n/2$ (this case must be dealt with separately). By symmetry we may assume that $k \geq n/2$. Then $0 < w \leq n$. If $w \geq n/2$ then the only solution is 0^n . If $w < n/2$, then let s be the unique integer $0 \leq s < \infty$, that satisfies $n/(12s + 14) \leq w < n/(12s + 2)$.

Case I: If $n/8 \leq w < n/2$ then the solutions are:

$$0^{\lfloor \frac{n}{2} \rfloor} 1^{n - \lfloor \frac{n}{2} \rfloor - w - c_1} 0^{w + c_1}$$

where $c_1 \in \{-1, 0, 1\}$.

Case II: If $n/(12s + 8) \leq w < n/(12s + 2)$ then the solutions are

$$\begin{cases} 0^{4w + c_1} 1^{\lfloor n/2 \rfloor - 4w - c_1} 0^{n - \lfloor n/2 \rfloor - 7w - (c_2 + c_3 + c_4)} 1^{6w + c_3} 0^{w + c_4} & \text{for } s = 1; \\ 0^{4w + c_4} (1^{6w + c_5^{s_i}} 0^{6w + c_6^{s_i}})^{s/2} Q(0^{6w + c_7^{s_i}} 1^{6w + c_8^{s_i}})^{s/2} 0^{w + c_9} & \text{for } s > 1. \end{cases}$$

where the c_j 's and $c_j^{s_i}$'s are bounded constants (independent of n) and Q can be an (almost) arbitrary mix of r zeroes and ones (where r is the unique integer such that the length of this interval is n). Further, the number of ones in Q is at most $12w$. Notation: (1) the $c_j^{s_i}$'s can be different constants with i ranging from 1 to $s/2$; (2) if s is odd $(ab)^{s/2}$ is $(ab)^{(s-1)/2}a$.

Case III: If $n/(12s + 14) \leq w < n/(12s + 8)$ then the solutions are

$$\begin{cases} 0^{4w + d_1} 1^{n - 5w - (d_1 + d_2)} 0^{w + d_2} & \text{for } s = 0; \\ 0^{4w + d_3} (1^{6w + d_4^{s_i}} 0^{6w + d_5^{s_i}})^{s/2} Q(0^{6w + d_6^{s_i}} 1^{6w + d_7^{s_i}})^{s/2} 0^{w + d_8} & \text{for } s > 0. \end{cases}$$

where the d_j 's and $d_j^{s_i}$'s are bounded constants (independent of n)

and Q can be an (almost) arbitrary mix of r zeroes and ones, with the number of ones in Q at most $6w$.

Case IV: if $w = 0$ (i.e. $s = \infty$), the solutions are:

$$0^{g_1} (1^{g_2^{n_i}} 0^{g_3^{n_i}})^{n/(2G_1)} Q(0^{g_4^{n_i}} 1^{g_5^{n_i}})^{n/(2G_2)}$$

where $g_1 \in \{0, 1, 2\}$, the other g_i 's and $g_i^{n_i}$'s are bounded between 3 and 11, Q is an (almost) arbitrary mix of r zeroes and ones with the number of ones bounded between 0 and 22, $G_1 = \sum_i (g_2^{n_i} + g_3^{n_i})$, and $G_2 = \sum_i (g_4^{n_i} + g_5^{n_i})$.

Proof: Routine verification!

Now it is time to impose the extra condition that $\sum_{i=1}^n z_i = k$ ($= (w + n)/2$). With Cases I and II a routine calculation yields a contradiction of the applicable range of w when n is sufficiently large. For Case III, a routine calculation yields a local minimum of $w = n/11$ if $s = 0$. If $s > 0$ argue as follows. Let t be the number of 1's in Q . Recall that r is the total number of 0's and

1's in Q . Let $w_c(s) = n/(12s + c)$ where we must have $8 \leq c \leq 14$. Since we need $\sum_{i=1}^n z_i = k$ ($= (w+n)/2$), we see that $6w_c(s)s + t = n(12s + c + 1)/(24s + 2c)$ gives $t = (c+1)w_c(s)/2$. Further, since the number of 1's in Q is bounded by $6w_c(s)$, we find that we must have $c \leq 11$. We also must have $r = n - w_c(s)(12s + 5)$, by the definition of r . Using the simple inequality $r \geq t$, we have $n - w_c(s)(12s + 5) \geq (c+1)w_c(s)/2$. From this deduce that $c \geq 11$. Hence we must have $c = 11$ at a local minimum. Thus the local minimums for Case III, $s > 0$, are $w_s = n/(12s + 11)$. Case IV gives infinitely many local minimums. Hence

The Local Minima Are Asymptotically Equivalent (mod $O(n)$) to:

$$\begin{cases} Z_s := 0^{4w_s} (1^{6w_s} 0^{6w_s})^{\frac{s}{2}} 1^{6w_s} (0^{6w_s} 1^{6w_s})^{\frac{s}{2}} 0^{w_s} & \text{for } 0 \leq s < \infty \text{ (where } w_s := \frac{n}{12s+11}), \\ Z_\infty^t = (0^t 1^t)^{n/(2t)} & \text{for } 3 \leq t \leq 11 \end{cases}$$

A routine calculation [R] shows that for $0 \leq s < \infty$

$$F(Z_s) = \frac{12s + 8}{16(12s + 11)} n^2 + O(n),$$

which is strictly increasing in s . An easy calculation shows $F(Z_\infty^t) = (1/16)n^2 + O(n)$ for any natural number t .

...And The Winner Is: $Z_0 = 0^{4n/11} 1^{6n/11} 0^{n/11}$ setting the world-record of $(1/22)n^2 + O(n)$.

An Extension Here we note that our result implies a good upper bound for the general r -coloring of the first n integers; if we r -color the integers (with colors $C_1 \dots C_r$) from 1 to n then the minimum number of monochromatic Schur triples is bounded above by

$$\frac{n^2}{2^{2r-3} 11} + O(n).$$

This comes from the following coloring:

$$\begin{cases} Color(i) = C_j & \text{if } \frac{n}{2^j} < i \leq \frac{n}{2^{j-1}} \quad \text{for } 1 \leq j \leq r-2, \\ Color(i) = C_{r-1} & \text{if } 1 \leq i \leq \frac{4n}{2^{r-2} 11} \text{ or } \frac{10n}{2^{r-2} 11} < i \leq \frac{n}{2^{r-2}}, \\ Color(i) = C_r & \text{if } \frac{4n}{2^{r-2} 11} < i \leq \frac{10n}{2^{r-2} 11}. \end{cases}$$

Note: Tomasz Schoen[S], a student of Tomasz Luczak, has independently solved this problem.

REFERENCES

[GRR] R. Graham, V. Rödl, and A. Rucinski, *On Schur properties of random subsets of integers*, J. Number Theory **61** (1996), 388-408.

[RZ] A. Robertson, *On the asymptotic behavior of Schur triples*, [www.math.temple.edu/~aaron].

[S] T. Schoen, *On the number of monochromatic Schur triples*, in preparation, [wtg3@informatik.uni-kiel.de].