



ON THE SUM OF RECIPROCAL OF AMICABLE NUMBERS

Jonathan Bayless*Department of Mathematics, Husson University, Bangor, Maine*
BaylessJ@husson.edu**Dominic Klyve***Department of Mathematics, Central Washington University, Ellensburg,*
Washington
klyved@cwu.edu*Received: 2/1/10, Revised: 9/24/10, Accepted: 12/26/10, Published: 3/9/11***Abstract**

Two numbers m and n are considered amicable if the sum of their proper divisors, $s(n)$ and $s(m)$, satisfy $s(n) = m$ and $s(m) = n$. In 1981, Pomerance showed that the sum of the reciprocals of all such numbers, P , is a constant. We obtain both a lower and an upper bound on the value of P .

1. Introduction

Since at least the time of the Ancient Greeks, amicable numbers have enjoyed the attention of mathematicians. Let $s(n)$ denote the sum of the proper divisors of n , that is, $s(n) = \sigma(n) - n$, where $\sigma(n)$ denotes Euler's sum-of-divisors function. Then an *amicable pair* is a pair of distinct integers m, n , such that $s(n) = m$ and $s(m) = n$. We will also refer to any integer which is a member of an amicable pair as an *amicable number*. The smallest amicable pair, $(220, 284)$, was known to Pythagoras c. 500 BCE. The study of amicable pairs was a topic arising often in Medieval Islam; as early as the 9th Century, Thābit had discovered three pairs, including $(17296, 18416)$ – a pair which was rediscovered independently by Borho, Ibn al-Bannā', Kamaladdin Fārisī, and Pierre de Fermat [4]. In the 18th century, Euler [1] famously advanced the theory of amicable numbers by giving a table of 30 new pairs in 1747 (one of which is, in fact, an error – see [13]), and a larger table of 61 pairs, together with a method of generating them, in 1750 [2].

Today, although much is known about amicable numbers (and their less popular friends, sociable numbers), there is still a lot that we don't know. For example,

although the best upper bound on their density was given by Pomerance [11] in 1981 (see (2) below), no known lower bound exists – indeed, it has not been proven that there are infinitely many. Considerable work has been done on questions such as the properties of even-even pairs, odd-odd pairs, pairs for which each number has exactly two prime factors not contained in its pair, and many more complex variations. For a nice survey on amicable numbers, see [4]. One interesting fact, which motivated the present work, is that the sum of the reciprocals of the amicable numbers converges, that is,

$$\sum_{a \text{ amicable}} \frac{1}{a} = P < \infty. \tag{1}$$

This is a consequence of the bound shown 28 years ago by Pomerance [11], but to date no bounds have been given on the value of this sum. In this paper, we provide an upper bound and a lower bound for the value of this sum.

1.1. The Distribution of Amicable Numbers

Following [11], we define $A(x)$ to be the count of amicable numbers not greater than x . In 1955, Erdős [3] showed that the amicable numbers have density zero. Pomerance showed [10]

$$A(x) \leq x \exp\left(-c\sqrt{\log \log \log x \log \log \log x}\right)$$

for a positive constant c and all sufficiently large x . Then, in 1981, Pomerance [11] improved this result to

$$A(x) \leq C \frac{x}{\exp\left((\log x)^{1/3}\right)} \tag{2}$$

for some constant C and all sufficiently large x . From this and partial summation, it is clear that we have (1). Since Pomerance was the first to show this sum to converge, we refer to its value, P , as *Pomerance’s constant*.

Having established that P is finite, it is natural to ask about its value. We prove the following theorem.

Theorem 1. *Pomerance’s constant, P , the sum of the the reciprocals of the amicable numbers, can be bounded as*

$$.0119841556 < P < 6.56 \times 10^8.$$

It should be noted that the upper bound in this result can be improved by a more careful choice of the functions in Table 1 on page 3. It is unclear how to fully optimize this argument, but the authors have been able to show $P < 3.4 \times 10^6$.

To establish Theorem 1, we split the sum defining P into three parts:

$$\sum_{\text{amicable } n} \frac{1}{n} = \sum_{\substack{\text{amicable } n \\ n \leq 10^{14}}} \frac{1}{n} + \sum_{\substack{\text{amicable } n \\ 10^{14} < n \leq \exp(10^6)}} \frac{1}{n} + \sum_{\substack{\text{amicable } n \\ n > \exp(10^6)}} \frac{1}{n}. \tag{3}$$

We evaluate the first sum directly in Section 2 to establish the lower bound for P . In Section 3, we bound the second sum. We then modify the argument in [11] to address the final sum above in Section 5.

Throughout, p , q , and r will denote primes, $P(n)$ the largest prime factor of n , and $\psi(x, y)$ the number of y -smooth numbers up to x , that is, the size of the set

$$\{n \leq x \mid P(n) \leq y\}.$$

Finally, a and b will represent amicable numbers.

We will repeatedly make use of the following functions. These will be redefined at the appropriate place, but the reader can use this table as a helpful reference.

x_0	$= \exp(10^6)$
y_0	$= \exp(26000)$
c	$= c(x) = 1 - (\log x)^{-1/6} (\log \log x)^{-1}$
c_0	$= c(10^6) > 0.99276$
σ	$= \sigma(y) = 1 - 1/(2 \log y)$
ℓ	$= \ell(x) = \exp\left((\log x)^{1/6}\right)$
L	$= L(x) = \exp\left(0.1882 (\log x)^{2/3} \log \log x\right)$

Table 1: A table of functions used in the proof of Theorem 1.

2. A Lower Bound – Some Reciprocal Sums

Determining a lower bound on P is straightforward – we need only sum the reciprocals of any subset of the set of amicable numbers to find one. Let P_j be the sum of all integers not greater than 10^j which are members of an amicable pair, i.e.,

$$P_j = \sum_{\substack{a \leq 10^j \\ a \text{ amicable}}} \frac{1}{a}.$$

The current record for exhaustive searches for amicable numbers is 10^{14} [9]. In Table 2, we give values for P_j for various powers of 10 up to this bound.

j	P_j
1	0
2	0
3	0.0080665813060179257
4	0.0111577261442474466
5	0.0117423756996823562
6	0.0119304720866743157
7	0.0119714208511438135
8	0.0119812212551025145
9	0.0119834313702743716
10	0.0119839922963130553
11	0.0119841199294457703
12	0.0119841486963721084
13	0.0119841542458770555
14	0.0119841556796931142

Table 2: Sums of reciprocals of amicable numbers to 10^j .

From this table, we immediately have

$$P > P_{14} > .0119841556796931142. \tag{4}$$

We note that this series seems to converge rather quickly. Indeed, for any $j \leq 13$ with $P_j > 0$, the difference between P_{14} and P_j is less than the difference between P_j and P_{j-1} . With this observation, we conjecture that the true value of P can be estimated by

$$|P - P_{14}| < P_{14} - P_{13} \approx .0000000006338,$$

and thus:

Conjecture 2. Pomerance’s constant, P , satisfies $P < .0119841563134$.

This, however, is merely conjecture. We turn now to establish the upper bound of Theorem 1.

3. The Middle Range

The amicable numbers in the range $[10^{14}, \exp(10^6)]$ are too large to be found explicitly, and too small to be amenable to the results we make use of later in this paper. While there are some ways to restrict the sum over these numbers, their final contribution to Pomerance’s constant is small, and we here simply make use of the trivial bound:

$$\sum_{10^{14} < n \leq \exp(10^6)} \frac{1}{n} < 10^6.$$

4. Preliminaries

We will use a number of explicit formulas of prime functions. For instance, we will use the fact (see [12, Theorem 5]) that

$$\sum_{p \leq x} \frac{1}{p} \leq \log \log x + B + \frac{1}{2(\log x)^2}$$

for $x \geq 286$, where $B = .26149721 \dots$. Recall that we have chosen $x_0 = \exp(10^6)$. This gives that, for $x \geq x_0$,

$$\sum_{p \leq x} \frac{1}{p} \leq \log \log x + .2615. \tag{5}$$

We will use a bound on the sum of reciprocals of primes in a particular residue class modulo p , where $p > 10^{14}$.

Lemma 3. *For any value of $y \geq 10^{14}$ and any prime $p \geq 10^{14}$, the following holds uniformly:*

$$\sum_{\substack{q \leq y \\ q \equiv -1 \pmod p}} \frac{1}{q} \leq \frac{4 + 3 \log \log y}{p}.$$

Proof. We begin with the Brun-Titchmarsh inequality of [8]; namely, for coprime integers k and n , the number of primes $q \leq y$ with $q \equiv k \pmod n$, denoted $\pi(y; n, k)$, satisfies

$$\pi(y; n, k) \leq \frac{2y}{\varphi(n) \log(y/n)} \text{ for } y > n.$$

For prime $p \geq 10^{14}$ and $(k, p) = 1$, we can use the fact that $\varphi(p) = p - 1$ to see that

$$\pi(y; p, k) \leq \frac{2y}{\varphi(p) \log(y/p)} \leq \frac{2.0001y}{p \log(y/p)}. \tag{6}$$

Note that the first prime $q \equiv -1 \pmod p$ is at least $2p - 1$, so we need only consider $p \leq \frac{y+1}{2}$. Thus, $\frac{y}{p} \geq 2 - \frac{1}{p} \geq 1.999999$ for $p \geq 10^{14}$. We use partial summation to

obtain

$$\begin{aligned} \sum_{\substack{q \leq y \\ q \equiv -1 \pmod p}} \frac{1}{q} &= \frac{\pi(y; p, -1)}{y} + \int_{2p-1}^y \frac{\pi(t; p, -1)}{t^2} dt \\ &\leq \frac{2.0001}{p \log(y/p)} + \int_{2p-1}^y \frac{2.0001}{p t \log(t/p)} dt \\ &= \frac{2.0001}{p \log(y/p)} + \frac{2.0001}{p} \int_{2p-1}^y \frac{dt}{t \log(t/p)} \\ &= \frac{2.0001}{p \log(y/p)} + \frac{2.0001}{p} \cdot \log \log(t/p) \Big|_{2p-1}^y \\ &= \frac{2.0001}{p} \left(\frac{1}{\log(y/p)} + \log \log(y/p) - \log \log \left(2 - \frac{1}{p} \right) \right). \end{aligned}$$

Since $\frac{1}{\log(y/p)} - \log \log \left(2 - \frac{1}{p} \right) < 1.8093$ and $\log \log(y/p) \leq \log \log y$, we see that

$$\sum_{\substack{q \leq y \\ q \equiv -1 \pmod p}} \frac{1}{q} \leq \frac{4 + 3 \log \log y}{p},$$

which proves the lemma. □

We will need the following lemma to bound the number of amicable numbers up to x which are also y -smooth for some $y \geq y_0 = \exp(26000)$. Recall that $\sigma = 1 - 1/(2 \log y)$.

Lemma 4. *For $y \geq y_0$ and σ as above, we have*

$$\prod_{p \leq y} \left(1 - \frac{1}{p^\sigma} \right)^{-1} < 7.6515 \log y.$$

Before proving Lemma 4, we establish a few necessary lemmas.

Lemma 5. *For $y \geq y_0$,*

$$\sum_{p \leq y} \frac{\log p}{p} \leq \log y - 1.3325.$$

Proof. We recall a result from Rosser and Schoenfeld (see [12, 3.23]). Let $E = -\gamma - \sum_p (\log p)/p^n < -1.332582275$. For $y \geq 319$,

$$\sum_{p \leq y} \frac{\log p}{p} \leq \log y + E + \frac{1}{2 \log y}.$$

Setting $y = y_0$, Lemma 5 follows. □

Lemma 6. *Let $y \geq y_0$. Then, for σ as above,*

$$\sum_{p \leq y} \frac{1}{p^\sigma} < \log \log y + 1.0859.$$

Proof. We first note the helpful fact that

$$p^\sigma = p \exp(-\log p / (2 \log y)) \geq p \exp(-1/2). \tag{7}$$

The idea in this proof is to bound the difference between $1/p$ and $1/p^\sigma$. Let $f(y) = p^{-y}$; then we want to estimate $f(\sigma) - f(1)$. Since $f'(y) = -(\log p)f(y)$, it follows from the mean value theorem that $f(\sigma) - f(1) = (1 - \sigma)(\log p)/p^y$ for some y between σ and 1. Since $\sigma = 1 - 1/(2 \log y)$, this gives

$$\begin{aligned} f(\sigma) - f(1) &\leq \frac{1}{2 \log y} \left(\frac{\log p}{p^\sigma} \right) \\ &\leq \frac{e^{1/2}}{2 \log y} \left(\frac{\log p}{p} \right), \end{aligned}$$

by (7).

Hence, by Lemma 5, we have

$$\sum_{p \leq y} \left(\frac{1}{p^\sigma} - \frac{1}{p} \right) \leq \frac{e^{1/2}}{2 \log y} \sum_{p \leq y} \frac{\log p}{p} \leq \frac{e^{1/2}}{2} \left(\frac{\log y - 1.3325}{\log y} \right) < 0.8244.$$

So this means that

$$\sum_{p \leq y} \frac{1}{p^\sigma} \leq 1 + \sum_{p \leq y} \frac{1}{p} \leq \log \log y + 1.0859,$$

by (5). □

Lemma 7. *For $y \geq y_0$ and σ as above, we have*

$$\sum_p \frac{1}{p^\sigma(p^\sigma - 1)} < .7734. \tag{8}$$

Proof. Since $y \geq y_0$, it follows that $\sigma > .9999$. We will bound (8) by explicit computation and analytic methods. Using 500000 as a useful place to split our computation, we write

$$\begin{aligned} \sum_p \frac{1}{p^\sigma(p^\sigma - 1)} &\leq \sum_{p \leq 500000} \frac{1}{p^\sigma(p^\sigma - 1)} + \sum_{500000 < p < \infty} \frac{1}{p^\sigma(p^\sigma - 1)} \\ &\leq .7733545 + \int_{500000}^{\infty} \frac{1}{t^\sigma(t^\sigma - 1)} dt \\ &\leq .7733545 + .0000001 \\ &\leq .7734, \end{aligned}$$

by explicit computation. □

We may now prove Lemma 4.

Proof.

$$\begin{aligned} \prod_{p \leq y} \left(\frac{1}{1 - 1/p^\sigma} \right) &\leq \exp \left(\sum_{p \leq y} \left(\frac{1}{p^\sigma} + \frac{1}{p^{2\sigma}} + \dots \right) \right) \\ &= \exp \left(\sum_{p \leq y} \frac{1}{p^\sigma} + \frac{1}{p^\sigma(p^\sigma - 1)} \right) \\ &= \exp \left(\sum_{p \leq y} \frac{1}{p^\sigma} \right) \cdot \exp \left(\sum_{p \leq y} \frac{1}{p^\sigma(p^\sigma - 1)} \right) \\ &\leq \exp(\log \log y + 1.0859) \exp(.7734) \\ &\leq 6.4193 \log y, \end{aligned}$$

by Lemmas 7 and 6. □

We now show that our choice of $x \geq x_0$ is “sufficiently large” to adapt Pomerance’s argument [11]. Let

$$c = 1 - (\log x)^{-1/6} (\log \log x)^{-1}.$$

For $x \geq x_0$, it is clear that

$$c \geq c_0 = 1 - (10^6)^{-1/6} (\log 10^6)^{-1} > .99276.$$

Lemma 8. *For any $c \geq c_0$ above, we have*

$$\sum_{k \geq 2} \frac{1}{k^c (k^c - 1)} \leq 1.0225.$$

Proof. First, we split the sum into two parts, i.e.,

$$\sum_{k \geq 2} \frac{1}{k^c (k^c - 1)} = \sum_{2 \leq k < 10^6} \frac{1}{k^c (k^c - 1)} + \sum_{k \geq 10^6} \frac{1}{k^c (k^c - 1)}.$$

For any value of $k \geq 10^6$, we have $k^c < 1.000002(k^c - 1)$, so the second sum can be bounded by

$$\sum_{k \geq 10^6} \frac{1}{k^c (k^c - 1)} \leq 1.000002 \sum_{k \geq 10^6} \frac{1}{k^{2c}} \leq 1.000002 \int_{10^6-1}^{\infty} \frac{dt}{t^{2c}} \leq .0000013.$$

A quick computation shows that

$$\sum_{2 \leq k < 10^6} \frac{1}{k^c (k^c - 1)} \leq 1.02247315,$$

and adding these two together completes the proof of the lemma. □

Lemma 9. For any $c \geq c_0$, we have

$$\sum_{p \geq 2} \frac{1}{p^c (p^c - 1)} \leq .7877.$$

Proof. As before, we split the sum into two parts, i.e.,

$$\sum_{p \geq 2} \frac{1}{p^c (p^c - 1)} = \sum_{2 \leq p < 10^6} \frac{1}{p^c (p^c - 1)} + \sum_{p \geq 10^6} \frac{1}{p^c (p^c - 1)}.$$

Once again, we use an explicit computation to show that

$$\sum_{2 \leq p \leq 10^6} \frac{1}{p^c (p^c - 1)} \leq .7876817684,$$

and the second sum can be bounded by

$$\sum_{p \geq 10^6} \frac{1}{p^c (p^c - 1)} \leq .0000013$$

as in the previous lemma. Adding these two together completes the proof of the lemma. \square

In Section 5, we will use the function $\ell = \exp\left((\log x)^{1/6}\right)$, which is referenced in the following lemma.

Lemma 10. For any $x \geq x_0$ and $c = 1 - (\log \log x)^{-1} (\log x)^{-1/6}$, we have

$$\sum_{p \leq \ell^4} \frac{1}{p^c} \leq \frac{1}{6} \log \log x + 2.0346.$$

Proof. First, note that

$$\begin{aligned} p^c &= p^{1 - (\log \log x)^{-1} (\log x)^{-1/6}} = p \exp\left(-\frac{\log p}{(\log x)^{1/6} (\log \log x)}\right) \\ &\geq p \exp\left(-\frac{4 (\log x)^{1/6}}{(\log x)^{1/6} (\log \log x)}\right) \\ &= p \exp\left(-\frac{4}{(\log \log x)}\right) \\ &\geq .7486p, \end{aligned}$$

since $p \leq \ell^4$ and $x \geq e^{10^6}$. Then, note that the mean value theorem says that, with $f(y) = p^{-y}$ and $f'(y) = -(\log p)/p^y$,

$$f(c) - f(1) \leq \frac{(1 - c)(\log p)}{p^c}.$$

Applying both of these facts, we see that

$$\sum_{p \leq \ell^4} \left(\frac{1}{p^c} - \frac{1}{p} \right) \leq 1.3358(1-c) \sum_{p \leq \ell^4} \frac{\log p}{p}.$$

By Lemma 5, this can be bounded by

$$\begin{aligned} (1-c)1.3358(\log \ell^4 - 1.3325) &= \frac{1.3358(4(\log x)^{1/6} - 1.3325)}{(\log x)^{1/6}(\log \log x)} \\ &\leq \frac{5.3432}{\log \log x} \\ &\leq .3868. \end{aligned}$$

Then, we have

$$\begin{aligned} \sum_{p \leq \ell^4} \frac{1}{p^c} &= \sum_{p \leq \ell^4} \frac{1}{p} + \sum_{p \leq \ell^4} \left(\frac{1}{p^c} - \frac{1}{p} \right) \\ &\leq \log \log \ell^4 + .3868 + .2615 \\ &\leq \frac{1}{6} \log \log x + 2.0346. \end{aligned}$$

by (5). □

In [11], Pomerance uses a function $N(s)$ which, for

$$\psi(m) = \prod_{q^a \parallel m} (q+1)q^{a-1},$$

is the number of m with $\psi(m) \leq s$ and $P(\psi(m))$ bounded by a somewhat small function of x . His bound applies only to “large x ”, so we must verify that our choice of $x \geq x_0$ is large enough. Pomerance showed that

$$N(s) \leq 2s^c \prod_{\substack{P(k) < \ell^4 \\ k > 1}} (1 - k^{-c})^{-1}.$$

We now follow [11], making the bounds explicit as we proceed.

$$\prod_{\substack{P(k) < \ell^4 \\ k > 1}} (1 - k^{-c})^{-1} = \prod_{\substack{P(k) < \ell^4 \\ k > 1}} \left(1 + \frac{1}{k^c} + \frac{1}{k^{2c}} + \dots \right). \tag{9}$$

Since $1 + t \leq e^t$, we have (for $t = \frac{1}{k^c} + \frac{1}{k^{2c}} + \dots$)

$$1 + \frac{1}{k^c} + \frac{1}{k^{2c}} + \dots \leq \exp \left(\frac{1}{k^c} + \frac{1}{k^{2c}} + \dots \right).$$

Putting this into (9) gives

$$\prod_{\substack{P(k) < \ell^4 \\ k > 1}} (1 - k^{-c})^{-1} \leq \exp \left(\sum_{\substack{P(k) < \ell^4 \\ k > 1}} k^{-c} + \sum_{\substack{P(k) < \ell^4 \\ k > 1}} (k^{-2c} + k^{-3c} + \dots) \right). \quad (10)$$

We may bound

$$\sum_{\substack{P(k) < \ell^4 \\ k > 1}} (k^{-2c} + k^{-3c} + \dots) \leq \sum_{k > 1} \frac{1}{k^c (k^c - 1)} \leq 1.0225,$$

by Lemma 8. Setting aside this secondary term, the remaining term in the exponential in (10) satisfies

$$\sum_{\substack{P(k) < \ell^4 \\ k > 1}} k^{-c} = \prod_{p < \ell^4} (1 - p^{-c})^{-1}.$$

Bounding this product by the same argument as in (9) and (10), we have

$$\begin{aligned} \prod_{p < \ell^4} (1 - p^{-c})^{-1} &\leq \exp \left(\sum_{p < \ell^4} p^{-c} + \sum_{p < \ell^4} \frac{1}{p^c (p^c - 1)} \right) \\ &\leq \exp(.7877) \exp \left(\sum_{p < \ell^4} p^{-c} \right), \end{aligned}$$

by Lemma 9. An application of Lemma 10 shows that this product is bounded by

$$\prod_{p < \ell^4} (1 - p^{-c})^{-1} \leq \exp \left(\frac{1}{6} \log \log x + 2.8223 \right) \leq \exp(2.8223) (\log x)^{1/6}.$$

Thus,

$$\prod_{\substack{P(k) < \ell^4 \\ k > 1}} (1 - k^{-c})^{-1} \leq \exp \left(1.0225 + \exp(2.8223) (\log x)^{1/6} \right).$$

We will need the bound in (9) to hold for

$$s \geq L = \exp \left(0.1882 (\log x)^{2/3} \log \log x \right).$$

If $s \geq L$, the argument above shows that the inequality

$$\begin{aligned} \frac{N(s)}{s} &\leq 2s^{c-1} \exp\left(1.0225 + \exp(2.8223) (\log x)^{1/6}\right) \\ &= 2 \exp\left((c-1) \log s + 1.0225 + \exp(2.8223) (\log x)^{1/6}\right) \\ &\leq 2 \exp\left(-.1882 (\log x)^{1/2} + 1.0225 + \exp(2.8223) (\log x)^{1/6}\right) \\ &\leq \frac{2e^{1.0225}}{\ell^2} \end{aligned}$$

holds for x such that

$$2 + \exp(2.8223) \leq .1882 (\log x)^{1/3}, \tag{11}$$

which, in turn, holds true for $x \geq x_0$.

Lastly, we will make use of the fact that for $k \geq 2$,

$$\sum_{n \geq k} \frac{1}{n^2} < \frac{1}{k-1}. \tag{12}$$

5. Large Amicable Numbers

5.1. Outline of the Proof

Pomerance’s proof of (2) rests on an argument that five different types of numbers do not contribute much to the sum defining P , with a careful count of amicable numbers among the remaining integers up to x . For the sake of brevity, we include only the statement of each assumption, and the resulting bound which must be made explicit to bound the value of P . We make a similar argument here, though we bound $A(x)$ by a much smaller function so that our bound will apply for smaller x . Following the notation of [11], we will use our previously defined functions

$$\ell = \exp\left((\log x)^{1/6}\right) \text{ and } L = \exp\left(.1882 (\log x)^{2/3} \log \log x\right). \tag{13}$$

Note that ℓ is smaller and L is slightly larger than their equivalents in [11]. It will also be useful to have $z = 2x \log \log x$.

Call an integer n *admissible* if it satisfies each of the five conditions given in [11]. We summarize them here.

- (i) The largest prime factor of n and $s(n)$ are both at least L^2 .
- (ii) If k^s divides n or $s(n)$ with $s \geq 2$, then $k^s < \ell^3$.
- (iii) If $p \mid \gcd(n, \sigma(n))$, then $p < \ell^4$.

(iv) The integer n satisfies

$$\frac{n}{P(n)} \geq L \quad \text{and} \quad \frac{s(n)}{P(s(n))} \geq L.$$

(v) If $m = \frac{n}{P(n)}$ and $m' = \frac{s(n)}{P(s(n))}$, then we have $P(\sigma(m)) \geq \ell^4$ and $P(\sigma(m')) \geq \ell^4$.

In showing (2), Pomerance proved the following (see [11] for details).

Theorem 11 (Pomerance). *The set of integers $n \leq x$ which are not admissible is $O\left(x/\exp\left((\log x)^{1/3}\right)\right)$. Specifically, the number of integers $n \leq x$ failing conditions (i)-(v) above are bounded, respectively, by the functions*

- (i) $\psi(z, L^2)$,
- (ii) $z \sum_{k^s \geq \ell^3} k^{-s}$,
- (iii) $\sum_{p \geq \ell^4} \sum_{\substack{q \equiv -1(p) \\ q \leq x/p}} \frac{x}{pq}$,
- (iv) $\frac{4x \log \log x}{L}$, and
- (v) $\frac{4e^{1.0225} x \log \log x (1 + 2 \log \log x)}{\ell^2} \sum_{p \leq x/L} \frac{1}{p}$.

Unfortunately, in [11], the bounds in (v) are only valid for very large x . By choosing the values of ℓ and L in (13), which are much smaller and slightly larger, respectively, than their counterparts in [11], we can make the argument apply for all $x \geq x_0 = \exp(10^6)$. In making these bounds explicit, we bound the quantity in (i) by the function C_1x/ℓ , the function in (ii) by C_2x/ℓ , and so on.

Let S denote the set of all admissible integers. Pomerance demonstrated that the count of amicable numbers in S can be bounded above by

$$\sum_{\substack{r \geq \ell^4 \\ q \leq x}} \sum_{q \equiv -1(r)} \sum_{\substack{m \equiv 0(q) \\ m \leq x}} \sum_{\substack{q' \equiv -1(r) \\ q' \leq z}} \frac{2x \log \log x}{q'm} \leq \frac{C_6x}{\ell} \tag{14}$$

for some constant C_6 (here (n) should be taken to mean mod n).

Then, it is clear that we may take $C = C_1 + C_2 + \dots + C_6$ in the bound

$$A(x) \leq C \frac{x}{\exp\left((\log x)^{1/6}\right)} \tag{15}$$

for $x \geq x_0$, and so it remains to find explicit values for each of these constants.

5.2. Evaluating the Constants

We address each of the constants given in Section 5.1.

C_1 : We use an explicit version of the method of Rankin to bound $\psi(z, L^2)$. Given that $x \geq x_0$, we also have that $L^2 \geq y_0$. Rankin’s method is based on the observation that, for any $\sigma > 0, x \geq 1$, and $y \geq 2$,

$$\begin{aligned} \psi(x, y) &\leq \sum_{\substack{n \geq 1 \\ P(n) \leq y}} \left(\frac{x}{n}\right)^\sigma = x^\sigma \sum_{\substack{n \geq 1 \\ P(n) \leq y}} \frac{1}{n^\sigma} \\ &= x^\sigma \prod_{p \leq y} \left(1 - \frac{1}{p^\sigma}\right)^{-1}. \end{aligned} \tag{16}$$

From Lemma 4 it follows that

$$\psi(x, y) \leq 6.4193 (x^\sigma) (\log y),$$

and, therefore, for $\sigma = 1 - 1/(2 \log y)$, that

$$\begin{aligned} \psi(z, L^2) &\leq 6.4193 \cdot z^\sigma \cdot .3764 (\log x)^{2/3} \log \log x \\ &\leq 2.4163 \cdot x^\sigma \cdot 2^\sigma (\log \log x)^{1+\sigma} \cdot (\log x)^{2/3}. \end{aligned}$$

Thus, we want to find C_1 such that

$$\frac{4.8325 \cdot x \cdot (\log \log x)^2 \cdot (\log x)^{2/3}}{z^{1/(.7528(\log x)^{2/3} \log \log x)}} \leq C_1 \frac{x}{\ell}.$$

Plugging in x_0 , we may take $C_1 = 13553617.97$.

It turns out that the other constants are negligible compared to C_1 , so we will sacrifice some sharpness in bounds for space in what follows.

C_2 : Note that we can bound

$$\begin{aligned} z \sum_{\substack{k^m \geq \ell^3 \\ m \geq 2}} \frac{1}{k^m} &< z \sum_{k \geq \ell^{\frac{3}{2}}} \left(\frac{1}{k^2} + \frac{1}{k^3} + \dots\right) = z \sum_{k \geq \ell^{\frac{3}{2}}} \frac{1}{k^2 - k} \\ &= z \sum_{k \geq \ell^{\frac{3}{2}}} \frac{1}{k(k-1)} < 1.0001z \sum_{k \geq \ell^{\frac{3}{2}}} \frac{1}{k^2} < \frac{1.0001z}{\ell^{\frac{3}{2}} - 1} \end{aligned}$$

for $x \geq x_0$. Now we need

$$\frac{1.0001z}{\ell^{\frac{3}{2}} - 1} < C_2 \frac{x}{\ell}.$$

Plugging in $x = x_0$ into the formula for ℓ shows that we may choose $C_2 = .1862$.

C_3 : A simple application of Lemma 3 gives

$$\frac{x}{p} \sum_{\substack{q \equiv -1(p) \\ q \leq x/p}} \frac{1}{q} \leq \frac{x}{p} \left(\frac{4 + 3 \log \log (x/p)}{p} \right).$$

Now, summing this over $p \geq \ell^4$ gives at most

$$x(4 + 3 \log \log x) \sum_{p \geq \ell^4} \frac{1}{p^2} < \frac{x(4 + 3 \log \log x)}{\ell^4 - 1},$$

by (12). Therefore, we need

$$\frac{x(4 + 3 \log \log x)}{\ell^4 - 1} \leq \frac{C_3 x}{\ell},$$

which holds for $C_3 > 4.3 \times 10^{-12}$.

C_4 : Here, we need

$$\frac{4x \log \log x}{L^2} < \frac{C_4 x}{\ell}$$

or

$$\frac{4 \log \log x \exp \left((\log x)^{1/6} \right)}{\exp \left(.61 (\log x)^{2/3} \log \log x \right)} < C_4.$$

Plugging in $x = x_0$ gives that we may choose any $C_4 > 10^{-17471}$.

C_5 : We must next compute C_5 such that the inequality

$$\frac{4e^{1.0225} x \log \log x (1 + 2 \log \log x)}{\ell^2} \sum_{p \leq x/\ell} \frac{1}{p} \leq \frac{C_5 x}{\ell}$$

holds for all $x \geq x_0$. By (5), this requires

$$\frac{4e^{1.0225} \log \log x (1 + 2 \log \log x)}{\exp \left((\log x)^{1/6} \right)} (\log \log x + .2615) < C_5.$$

For $x \geq x_0$, we may take $C_5 = 2.8117$.

C_6 : We have

$$\sum_{\substack{r \geq \ell^4 \\ q \equiv -1(r) \\ q \leq x}} \sum_{\substack{m \equiv 0(q) \\ m \leq x}} \sum_{\substack{q' \equiv -1(r) \\ q' \leq z}} \frac{2x \log \log x}{q'm} = z \sum_r \sum_q \sum_m \frac{1}{m} \sum_{q'} \frac{1}{q'}. \tag{17}$$

Applying Lemma 3, we find (17) is bounded by

$$z(4 + 3 \log \log z) \sum_r \sum_q \sum_m \frac{1}{m}. \tag{18}$$

Now, since $\log(x/q) < \log x$, we can bound

$$\sum_{\substack{m \equiv 0(q) \\ m \leq x}} \frac{1}{m} = \sum_{\substack{n \leq x/q \\ nq \leq x}} \frac{1}{nq} = \frac{1}{q} \sum_{n \leq x/q} \frac{1}{n} \leq \frac{\log x}{q}.$$

Using this and another application of Lemma 3, the iterated sum in (18) above can be bounded by

$$z(4 + 3 \log \log z)(\log x)(4 + 3 \log \log x) \sum_r \frac{1}{r^2}. \tag{19}$$

Lastly, (12) shows that

$$\sum_{r \geq \ell^4} \frac{1}{r^2} < \frac{1}{\ell^4 - 1},$$

and so, putting this into (19), we need

$$\frac{z(4 + 3 \log \log z)(\log x)(4 + 3 \log \log x)}{\ell^4 - 1} \leq C_6 \frac{x}{\ell}. \tag{20}$$

A calculation shows that (20) holds for $C_6 = .0054$.

Thus, for $x \geq \exp(10^6)$, the number of amicable numbers not greater than x , $A(x)$, can be bounded as

$$A(x) \leq 13553620.97 \frac{x}{\exp((\log x)^{1/6})}.$$

Then, by partial summation, we have that the third sum in (3) is bounded by

$$13553620.97 (\exp(-10) + 1) \int_{x_0}^{\infty} \frac{dt}{t \exp((\log t)^{1/6})} \leq 654666169.$$

Combined with the results of Sections 2 and 3, this proves Theorem 1.

Acknowledgements The authors would like to thank Carl Pomerance for many helpful suggestions, as well as for his expert guidance over the years. Thanks to Paul Pollack for his help in making some of the bounds in [11] explicit in Section 4, and Kevin Ford for a number of helpful ideas. We also thank an anonymous referee for many suggested improvements in the writing of this paper.

References

- [1] L. Euler, ‘De numeris amicabilebus’ (E100), *Nova acta eruditorum*, **1747**, 267–269, reprinted in the Opera Omnia, Series I volume 2, 59–61. Also available on line, with an English translation by Jordan Bell, at <http://www.eulerarchive.org/>.
- [2] L. Euler, ‘De numeris amicabilebus’ (E152), *Opuscula varii argumenti* **2**, **1750**, 23–107, reprinted in the Opera Omnia, Series I volume 2, 86–162. Also available on line at <http://www.eulerarchive.org/>.
- [3] P. Erdős, ‘On amicable numbers’, *Publ. Math. Debrecen* **4** (1955), 108–111.
- [4] M. Garca, J.M. Pedersen, and H.J.J. te Riele, ‘Amicable pairs, a survey’, *High Primes and Misdemeanours: Lectures in Honour of the 60th Birthday of Hugh Cowie Williams*, in: *Fields Inst. Commun.*, vol. **41**, Amer. Math. Soc., Providence, RI, 2004, 179–196.
- [5] H.-J. Kanold. ‘Über befreundete Zahlen II’, *Math. Nachr.*, **10** (1953), 99–111.
- [6] M. Kobayashi, P. Pollack, C. Pomerance. ‘On the distribution of sociable numbers.’ *Journal of Number Theory*, **129** (2009), 1990–2009.
- [7] E.J. Lee, ‘On the divisibility by nine of the sums of even amicable pairs’, *Math. Comp.* **23** (1969), 545–548.
- [8] H. Montgomery and R. Vaughan, ‘The large sieve’, *Mathematika* (20) **40** (1973), 119–134.
- [9] J. Pedersen, ‘Known Amicable Pairs’, electronic resource available at <http://amicable.homepage.dk/>.
- [10] C. Pomerance, ‘On the distribution of amicable numbers.’, *Journal fr die reine und angewandte Mathematik* **293/294** (1977), 217–222.
- [11] C. Pomerance, ‘On the distribution of amicable numbers. II’, *Journal fr die reine und angewandte Mathematik* **325** (1981), 183–188.
- [12] J. B. Rosser and L. Schoenfeld, ‘Approximate formulas for some functions of prime numbers’, *Illinois Journal of Mathematics* **6** (1962), 64–94.
- [13] E. Sandifer, ‘Amicable Numbers’, *How Euler Did It*, Math. Assoc. Amer. (2007), 49–56.