



## THE $(r_1, \dots, r_p)$ -BELL POLYNOMIALS

Mohammed Said Maamra

*Faculty of Mathematics, RECITS's Laboratory, USTHB, Algiers, Algeria*  
mmaamra@usthb.dz or mmaamra@yahoo.fr

Miloud Mihoubi

*Faculty of Mathematics, RECITS's laboratory,*  
*USTHB, Algiers, Algeria.*  
mmihoubi@usthb.dz or miloudmihoubi@gmail.com

*Received: 12/10/12, Revised: 12/21/13, Accepted: 4/19/14, Published: 7/10/14*

### Abstract

In a previous paper, Mihoubi et al. introduced the  $(r_1, \dots, r_p)$ -Stirling numbers and the  $(r_1, \dots, r_p)$ -Bell polynomials and gave some of their combinatorial and algebraic properties. These numbers and polynomials generalize, respectively, the  $r$ -Stirling numbers of the second kind introduced by Broder and the  $r$ -Bell polynomials introduced by Mező. In this paper, we prove that the  $(r_1, \dots, r_p)$ -Stirling numbers of the second kind are log-concave. We also give generating functions and generalized recurrences related to the  $(r_1, \dots, r_p)$ -Bell polynomials.

### 1. Introduction

In 1984, Broder [2] introduced and studied the  $r$ -Stirling number of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}_r$ , which counts the number of partitions of the set  $[n] = \{1, 2, \dots, n\}$  into  $k$  non-empty subsets such that the  $r$  first elements are in distinct subsets. In 2011, Mező [8] introduced and studied the  $r$ -Bell polynomials. In 2012, Mihoubi et al. [12] introduced and studied the  $(r_1, \dots, r_p)$ -Stirling number of the second kind  $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}_{r_1, \dots, r_p}$ , which counts the number of partitions of the set  $[n]$  into  $k$  non-empty subsets such that the elements of each of the  $p$  sets  $R_1 := \{1, \dots, r_1\}$ ,  $R_2 := \{r_1 + 1, \dots, r_1 + r_2\}$ ,  $\dots$ ,  $R_p := \{r_1 + \dots + r_{p-1} + 1, \dots, r_1 + \dots + r_p\}$  are in distinct subsets.

This work is motivated by the study of the  $r$ -Bell polynomials [8] and the  $(r_1, \dots, r_p)$ -Stirling numbers of the second kind [12], in which we may establish

- the log-concavity of the  $(r_1, \dots, r_p)$ -Stirling numbers of the second kind,
- generalized recurrences for the  $(r_1, \dots, r_p)$ -Bell polynomials, and
- the ordinary generating functions of these numbers and polynomials.

To begin, by the symmetry of the  $(r_1, \dots, r_p)$ -Stirling numbers with respect to  $r_1, \dots, r_p$ , let us suppose that  $r_1 \leq r_2 \leq \dots \leq r_p$ , and throughout this paper we use the following notation and definitions

$$\begin{aligned} \mathbf{r}_p &:= (r_1, \dots, r_p), \quad |\mathbf{r}_p| := r_1 + \dots + r_p, \\ P_t(z; \mathbf{r}_p) &:= (z + r_p)^t (z + r_p)^{r_1} \dots (z + r_p)^{r_{p-1}}, \quad t \in \mathbb{R}, \\ B_n(z; \mathbf{r}_p) &:= \sum_{k=0}^{n+|\mathbf{r}_{p-1}|} \left\{ \begin{matrix} n + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} z^k, \quad n \geq 0 \end{aligned}$$

and  $\mathbf{e}_i$  denotes the  $i$ -th vector of the canonical basis of  $\mathbb{R}^p$ . In [12], the following were proved:

$$B_n(z; \mathbf{r}_p) = \exp(-z) \sum_{k \geq 0} P_n(k; \mathbf{r}_p) \frac{z^k}{k!}, \tag{1}$$

$$P_n(z; \mathbf{r}_p) = \sum_{k=0}^{n+|\mathbf{r}_{p-1}|} \left\{ \begin{matrix} n + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} z^k. \tag{2}$$

For later use we define the following numbers

$$a_k(\mathbf{r}_{p-1}) = (-1)^{|\mathbf{r}_{p-1}|-k} \sum_{|\mathbf{j}_{p-1}|=k} \begin{bmatrix} r_1 \\ j_1 \end{bmatrix} \dots \begin{bmatrix} r_{p-1} \\ j_{p-1} \end{bmatrix}, \quad |\mathbf{j}_{p-1}| = j_1 + \dots + j_{p-1},$$

where  $\begin{bmatrix} n \\ k \end{bmatrix}$  are the absolute Stirling numbers of the first kind. Upon using the known identity

$$(u)^r = \sum_{j=0}^r (-1)^{r-j} \begin{bmatrix} r \\ j \end{bmatrix} u^j$$

we may state that we have

$$\sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) u^k = (u)^{r_1} \dots (u)^{r_{p-1}}. \tag{3}$$

In our contribution, we give more properties for the  $\mathbf{r}_p$ -Stirling numbers and the  $\mathbf{r}_p$ -Bell polynomials. The paper is organized as follows. In the next section we prove

that the sequence  $\left(\left\{\begin{matrix} n+|\mathbf{r}_p| \\ k+r_p \end{matrix}\right\}_{\mathbf{r}_p}; 0 \leq k \leq n + |\mathbf{r}_{p-1}|\right)$  is strongly log-concave and we give an approximation of  $\left\{\begin{matrix} n+|\mathbf{r}_p| \\ k+r_p \end{matrix}\right\}_{\mathbf{r}_p}$  when  $n \rightarrow \infty$  for a fixed  $k$ . In the third section we write  $B_n(z; \mathbf{r}_p)$  in the basis  $\{B_{n+k}(z; r_p) : 0 \leq k \leq |\mathbf{r}_{p-1}|\}$  and  $B_{n+m}(z; \mathbf{r}_p)$  in the family of bases  $\{z^j B_m(z; \mathbf{r}_p + j\mathbf{e}_p) : 0 \leq j \leq n\}$ . As consequences, we also give some identities for the  $\mathbf{r}_p$ -Stirling numbers. In the fourth section we give the ordinary generating functions of the  $\mathbf{r}_p$ -Stirling numbers of the second kind and the  $\mathbf{r}_p$ -Bell polynomials.

**2. Log-Concavity of the  $\mathbf{r}_p$ -Stirling Numbers**

In this section we discuss the real roots of the polynomial  $B_n(z; \mathbf{r}_p)$ , the log-concavity of the sequence  $\left(\left\{\begin{matrix} n+|\mathbf{r}_p| \\ k+r_p \end{matrix}\right\}_{\mathbf{r}_p}, 0 \leq k \leq n + |\mathbf{r}_{p-1}|\right)$ , the greatest maximizing index of  $\left\{\begin{matrix} n \\ k \end{matrix}\right\}_{\mathbf{r}_p}$  and we give an approximation of  $\left\{\begin{matrix} n+|\mathbf{r}_p| \\ m+r_p \end{matrix}\right\}_{\mathbf{r}_p}$  when  $n$  tends to infinity. The case  $p = 1$  was studied by Mező [9] and another study is done by Zhao [15] for a large class of the Stirling numbers.

In what follows, for illustration or if the order of  $r_1, \dots, r_p$  is unknown, we write the polynomial  $B_n(z; \mathbf{r}_p)$  as  $B_n(z; r_1, \dots, r_p)$  for which  $r_1, \dots, r_p$  are taken in any order.

**Theorem 1.** *The roots of the polynomial  $B_n(z; \mathbf{r}_p)$  are real and non-positive.*

To prove this theorem, we use the following lemma.

**Lemma 2.** *Let  $j, p$  be nonnegative integers and set*

$$B_n^{(j)}(z; \mathbf{r}_p) := \exp(-z) \frac{d^j}{dz^j} (z^{r_p} \exp(z) B_n(z; \mathbf{r}_p)),$$

where  $r_0 := 0$  and  $B_n(z; \mathbf{r}_0) := B_n(z) = \sum_{k=0}^n \left\{\begin{matrix} n \\ k \end{matrix}\right\} z^k$ . Then, we have

$$\begin{aligned} B_n^{(j)}(z; \mathbf{r}_p) &= z^{r_p-j} B_n(z; r_1, \dots, r_p, j) \quad \text{if } j < r_p, \\ B_n^{(j)}(z; \mathbf{r}_p) &= B_n(z; r_1, \dots, r_p, j) \quad \text{if } j \geq r_p, \end{aligned}$$

with  $\deg B_n^{(j)} = n + |\mathbf{r}_p|$ . In particular, we have  $B_n^{(r_{p+1})}(z; \mathbf{r}_p) = B_n(z; \mathbf{r}_{p+1})$ .

*Proof.* The definition of  $B_n^{(j)}(z; \mathbf{r}_p)$  and the identity (1) show that we have

$$\begin{aligned} & \exp(z) B_n^{(j)}(z; \mathbf{r}_p) \\ &= \frac{d^j}{dz^j} \left( \sum_{k \geq 0} P_n(k; \mathbf{r}_p) \frac{z^{k+r_p}}{k!} \right) \\ &= \sum_{k \geq \max(0, j-r_p)} (k+r_p)^n (k+r_p)^{r_1} \cdots (k+r_p)^{r_{p-1}} (k+r_p)^j \frac{z^{k+r_p-j}}{k!}. \end{aligned}$$

Then, for  $0 \leq j < r_p$  we obtain

$$\begin{aligned} \exp(z) B_n^{(j)}(z; \mathbf{r}_p) &= \sum_{k \geq 0} (k+r_p)^n (k+r_p)^{r_1} \cdots (k+r_p)^{r_{p-1}} (k+r_p)^j \frac{z^{k+r_p-j}}{k!} \\ &= z^{r_p-j} \exp(z) B_n(z; r_1, \dots, r_p, j) \end{aligned}$$

and for  $j \geq r_p$  we obtain

$$\begin{aligned} \exp(z) B_n^{(j)}(z; \mathbf{r}_p) &= \sum_{k \geq j-r_p} (k+r_p)^n (k+r_p)^{r_1} \cdots (k+r_p)^{r_{p-1}} (k+r_p)^j \frac{z^{k+r_p-j}}{k!} \\ &= \sum_{k \geq 0} (k+j)^n (k+j)^{r_1} \cdots (k+j)^{r_{p-1}} (k+j)^{r_p} \frac{z^k}{k!} \\ &= \exp(z) B_n(z; r_1, \dots, r_p, j). \end{aligned}$$

It is obvious that we have  $\deg B_n^{(j)} = n + |\mathbf{r}_p|$  and for  $j = r_{p+1} \geq r_p$  we obtain  $B_n^{(r_{p+1})}(z; \mathbf{r}_p) = B_n(z; r_1, \dots, r_p, r_{p+1}) = B_n(z; \mathbf{r}_{p+1})$ .  $\square$

*Proof of Theorem 1.* We will show by induction on  $p$  that the roots of the polynomials  $B_n(z; \mathbf{r}_p)$  are real and non-positive. Indeed, for  $p = 0$  the classical Bell polynomial  $B_n(z; \mathbf{r}_0) = B_n(z)$  has only real non-positive roots and for  $p = 1$  the polynomial  $B_n(z; \mathbf{r}_1)$  is the  $r_1$ -Bell polynomial introduced in [8] and has only real non-positive roots. Assume, for  $1 \leq r_1 \leq r_2 \leq \dots \leq r_p$ , that the roots of the polynomial  $B_n(z; \mathbf{r}_p)$  are real and negative, denoted by  $z_1, \dots, z_{n+|\mathbf{r}_{p-1}|}$  with  $0 > z_1 \geq \dots \geq z_{n+|\mathbf{r}_{p-1}|}$ . We will prove that the polynomial  $B_n^{(j)}(z; \mathbf{r}_p)$  has only real non-positive roots and we conclude that the polynomial  $B_n(z; \mathbf{r}_{p+1}) = B_n^{(r_{p+1})}(z; \mathbf{r}_p)$  (see Lemma 2) has only real non-positive roots.

Firstly, we examine the polynomials  $B_n^{(j)}(z; \mathbf{r}_p)$  for  $j < r_p$ . Indeed, the above statements show that the function

$$f_n(z; \mathbf{r}_p) := \exp(z) B_n^{(0)}(z; \mathbf{r}_p) = z^{r_p} \exp(z) B_n(z; \mathbf{r}_p)$$

vanishes at  $z_0, z_1, \dots, z_{n+|\mathbf{r}_{p-1}|}$  with  $z_0 = 0 > z_1 \geq \dots \geq z_{n+|\mathbf{r}_{p-1}|}$  and  $z_0 = 0$  is of multiplicity  $r_p$ . Lemma 2 gives

$$\frac{d}{dz} (f_n(z; \mathbf{r}_p)) = \exp(z) B_n^{(1)}(z; \mathbf{r}_p) = z^{r_p-1} \exp(z) B_n(z; r_1, \dots, r_{p-1}, r_p, 1)$$

and by applying Rolle's theorem to the function  $f_n(z; \mathbf{r}_p)$  we conclude that its derivative  $\frac{d}{dz} (f_n(z; \mathbf{r}_p))$  vanishes at some points  $x_1, \dots, x_{n+|\mathbf{r}_{p-1}|}$  with  $0 > x_1 \geq z_1 \geq x_2 \geq \dots \geq x_{n+|\mathbf{r}_{p-1}|} \geq z_{n+|\mathbf{r}_{p-1}|}$ . Consequently, the polynomial  $B_n^{(1)}(z; \mathbf{r}_p)$  vanishes at  $x_1, \dots, x_{n+|\mathbf{r}_{p-1}|}$  and at  $x_0 = 0$  (with multiplicity  $r_p - 1$ ). The number of these roots is  $(n + |\mathbf{r}_{p-1}|) + (r_p - 1) = n + |\mathbf{r}_p| - 1$ . Because  $B_n^{(1)}(z; \mathbf{r}_p)$  is of degree  $n + |\mathbf{r}_p|$  (see Lemma 2), it must have exactly  $n + |\mathbf{r}_p|$  finite roots; the missing one, denoted by  $x_{n+|\mathbf{r}_{p-1}|+1}$ , cannot be complex. By the fact that the coefficients of  $z^k$  in  $B_n(z; r_1, \dots, r_{p-1}, r_p, 1)$  are positive, the root  $x_{n+|\mathbf{r}_{p-1}|+1}$  must be negative too. So, the polynomial  $B_n^{(1)}(z; \mathbf{r}_p)$  has  $n + |\mathbf{r}_{p-1}| + 1$  real negative roots and  $z = 0$  is a root with multiplicity  $r_p - 1$ . Similarly, we apply Rolle's theorem to the function  $\frac{d}{dz} (f_n(z; \mathbf{r}_p))$  to conclude that the polynomial  $B_n^{(2)}(z; \mathbf{r}_p)$  has  $n + |\mathbf{r}_{p-1}| + 2$  real negative roots and  $z = 0$  is a root with multiplicity  $r_p - 2$ , and so on. So, the polynomials  $B_n^{(0)}(z; \mathbf{r}_p), B_n^{(1)}(z; \mathbf{r}_p), \dots, B_n^{(r_p-1)}(z; \mathbf{r}_p)$  have only real non-positive roots.

Secondly, we examine the polynomials  $B_n^{(j)}(z; \mathbf{r}_p)$  for  $r_p \leq j \leq r_{p+1}$ . Indeed, we have  $B_n^{(r_p)}(0; \mathbf{r}_p) \neq 0$  and consider the function

$$\frac{d^{r_p-1}}{dz^{r_p-1}} f_n(z; \mathbf{r}_p) = \exp(z) B_n^{(r_p-1)}(z; \mathbf{r}_p) = z \exp(z) B_n(z; r_1, \dots, r_{p-1}, r_p, r_{p-1}).$$

As it is shown above, this function has  $n + |\mathbf{r}_p| - 1$  real negative roots and the root  $z = 0$ , then Rolle's theorem shows that its derivative

$$\frac{d^{r_p}}{dz^{r_p}} f_n(z; \mathbf{r}_p) = \exp(z) B_n^{(r_p)}(z; \mathbf{r}_p) = \exp(z) B_n(z; r_1, \dots, r_{p-1}, r_p, r_p)$$

has at least  $n + |\mathbf{r}_p| - 1$  real negative roots. This means that the polynomial

$$B_n^{(r_p)}(z; \mathbf{r}_p) = B_n(z; r_1, \dots, r_{p-1}, r_p, r_p)$$

has at least  $n + |\mathbf{r}_p| - 1$  real negative roots and because it is of degree  $n + |\mathbf{r}_p|$ , the missing one cannot be complex. By the fact that the coefficients of  $z^k$  in  $B_n(z; r_1, \dots, r_{p-1}, r_p, r_p)$  are positive, this root must be negative too. So, the polynomial  $B_n^{(r_p)}(z; \mathbf{r}_p)$  has  $n + |\mathbf{r}_p|$  real negative roots. Similarly, apply Rolle's theorem to  $\frac{d^{r_p}}{dz^{r_p}} f_n(z; \mathbf{r}_p)$  and conclude that the polynomial  $B_n^{(r_p+1)}(z; \mathbf{r}_p)$  has  $n + |\mathbf{r}_p|$  real negative roots and so on. So, the polynomials  $B_n^{(r_p)}(z; \mathbf{r}_p), \dots, B_n^{(r_{p+1})}(z; \mathbf{r}_p)$  vanish only at negative numbers. Then, the polynomial  $B_n(z; \mathbf{r}_{p+1}) = B_n^{(r_{p+1})}(z; \mathbf{r}_p)$  (see Lemma 2) has only real negative roots.  $\square$

Upon using Newton’s inequality [6, p. 52], which is given by

**Theorem 3.** (*Newton’s inequality*) Let  $a_0, a_1, \dots, a_n$  be real numbers. If all the zeros of the polynomial  $P(x) = \sum_{k=0}^n a_k x^k$  are real, then the coefficients of  $P$  satisfy

$$a_i^2 \geq \left(1 + \frac{1}{i}\right) \left(1 + \frac{1}{n-i}\right) a_{i+1} a_{i-1}, \quad 1 \leq i \leq n-1,$$

we may state that:

**Corollary 4.** The sequence  $\left\{ \binom{n+|\mathbf{r}_p|}{k+r_p}_{\mathbf{r}_p}, 0 \leq k \leq n + |\mathbf{r}_{p-1}| \right\}$  is strongly log-concave (and thus unimodal).

This property shows that the sequence  $(\binom{n}{k}_{\mathbf{r}_p}, 0 \leq k \leq n)$  admits an index  $K \in \{0, 1, \dots, n\}$  for which  $\binom{n}{K}_{\mathbf{r}_p}$  is the maximum of  $\binom{n}{k}_{\mathbf{r}_p}$ . An application of Darroch’s inequality [3] will help us to localize this index.

**Theorem 5.** (*Darroch’s inequality*) Let  $a_0, a_1, \dots, a_n$  be real numbers. If all the zeros of the polynomial  $P(x) = \sum_{k=0}^n a_k x^k$  are real and negative and  $P(1) > 0$ , then the value of  $k$  for which  $a_k$  is maximized is within one of  $P'(1)/P(1)$ .

The following corollary gives a small interval for this index.

**Corollary 6.** Let  $K_{n,\mathbf{r}_p}$  be the greatest maximizing index of  $\binom{n}{k}_{\mathbf{r}_p}$ . We have

$$\left| K_{n+|\mathbf{r}_p|,\mathbf{r}_p} - \left( \frac{B_{n+1}(1; \mathbf{r}_p)}{B_n(1; \mathbf{r}_p)} - (r_p + 1) \right) \right| < 1.$$

*Proof.* Since the sequence  $\left\{ \binom{n+|\mathbf{r}_p|}{k+r_p}_{\mathbf{r}_p} \right\}$  is strongly log-concave, there exists an index  $K_{n+|\mathbf{r}_p|,\mathbf{r}_p}$  for which  $\binom{n+|\mathbf{r}_p|}{r_p}_{\mathbf{r}_p} < \dots < \binom{n+|\mathbf{r}_p|}{K_{n+|\mathbf{r}_p|,\mathbf{r}_p}}_{\mathbf{r}_p} > \dots > \binom{n+|\mathbf{r}_p|}{n+r_p}_{\mathbf{r}_p}$ . Then, on applying Theorem 1 and Darroch’s theorem, we obtain

$$\left| K_{n+|\mathbf{r}_p|,\mathbf{r}_p} - \frac{\frac{d}{dz} B_n(z; \mathbf{r}_p) \Big|_{z=1}}{B_n(1; \mathbf{r}_p)} \right| < 1.$$

It remains to use the first identity given in [12, Corollary 12] by  $z \frac{d}{dz} (B_n(z; \mathbf{r}_p)) = B_{n+1}(z; \mathbf{r}_p) - (z + r_p) B_n(z; \mathbf{r}_p)$ . □

### 3. Generalized Recurrences and Consequences

In this section, different representations of the polynomial  $B_n(z; \mathbf{r}_p)$  in different bases or families of basis are given by Theorems 7 and 10. Indeed, a representation in the basis  $\{B_{n+k}(z; r_p) : 0 \leq k \leq n + |\mathbf{r}_{p-1}|\}$  is given by the following theorem.

**Theorem 7.** *We have*

$$B_n(z; \mathbf{r}_p) = \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) B_{n+k}(z; r_p),$$

$$B_n(z; \mathbf{r}_{p+q}) = \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) B_{n+k}(z; r_p, \dots, r_{p+q}).$$

*Proof.* Upon using the fact that  $(k + r_p)^{r_m} = \sum_{j=0}^{r_m} (-1)^{r_m-j} \begin{bmatrix} r_m \\ j \end{bmatrix} (k + r_p)^j$ , we get

$$\begin{aligned} B_n(z; \mathbf{r}_p) &= \exp(-z) \sum_{k \geq 0} P_n(k; \mathbf{r}_p) \frac{z^k}{k!} \\ &= \exp(-z) \sum_{k \geq 0} \sum_{j=0}^{r_m} (-1)^{r_m-j} \begin{bmatrix} r_m \\ j \end{bmatrix} \frac{P_0(k; \mathbf{r}_p)}{(k + r_p)^{r_m}} (k + r_p)^{n+j} \frac{z^k}{k!} \\ &= \sum_{j=0}^{r_m} (-1)^{r_m-j} \begin{bmatrix} r_m \\ j \end{bmatrix} B_{n+j}(z; \mathbf{r}_p - r_m \mathbf{e}_m), \quad m = 1, 2, \dots, p-1, \end{aligned}$$

and with the same process, we obtain

$$\begin{aligned} B_n(z; \mathbf{r}_p) &= \sum_{j_1=0}^{r_1} \dots \sum_{j_{p-1}=0}^{r_{p-1}} (-1)^{|\mathbf{r}_{p-1}| - |\mathbf{j}_{p-1}|} \begin{bmatrix} r_1 \\ j_1 \end{bmatrix} \dots \begin{bmatrix} r_{p-1} \\ j_{p-1} \end{bmatrix} B_{n+|\mathbf{j}_{p-1}|}(z; r_p) \\ &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} (-1)^{|\mathbf{r}_{p-1}| - k} B_{n+k}(z; r_p) \sum_{|\mathbf{j}_{p-1}|=k} \begin{bmatrix} r_1 \\ j_1 \end{bmatrix} \dots \begin{bmatrix} r_{p-1} \\ j_{p-1} \end{bmatrix} \\ &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) B_{n+k}(z; r_p). \end{aligned}$$

This implies the first identity of the theorem.

Now, from Lemma 2 we can write

$$\begin{aligned} \exp(-z) \frac{d^{r_{p+1}}}{dz^{r_{p+1}}} (z^{r_p} \exp(z) B_n(z; \mathbf{r}_p)) \\ = \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) \exp(-z) \frac{d^{r_{p+1}}}{dz^{r_{p+1}}} (z^{r_p} \exp(z) B_{n+k}(z; r_p)), \end{aligned}$$

which gives by utilizing Lemma 2:  $B_n(z; \mathbf{r}_{p+1}) = \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) B_{n+k}(z; r_p, r_{p+1})$ .  
 We can repeat this process  $q$  times to obtain the second identity of the theorem.  $\square$

So, the  $\mathbf{r}_p$ -Stirling numbers admit an expression in terms of the usual  $r$ -Stirling numbers given by the following corollary.

**Corollary 8.** *We have*

$$\left\{ \begin{matrix} n + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} = \sum_{j=0}^{|\mathbf{r}_{p-1}|} \left\{ \begin{matrix} n + j + r_p \\ k + r_p \end{matrix} \right\}_{r_p} a_j(\mathbf{r}_{p-1}).$$

*Proof.* Using Theorem 7, the polynomial  $B_n(z; \mathbf{r}_p)$  can be written as follows:

$$\begin{aligned} \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) B_{n+j}(z; r_p) &= \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) \sum_{k=0}^{n+j} \left\{ \begin{matrix} n + j + r_p \\ k + r_p \end{matrix} \right\}_{r_p} z^k \\ &= \sum_{k=0}^{n+|\mathbf{r}_{p-1}|} z^k \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) \left\{ \begin{matrix} n + j + r_p \\ k + r_p \end{matrix} \right\}_{r_p} \end{aligned}$$

and since  $B_n(z; \mathbf{r}_p) = \sum_{k=0}^{n+|\mathbf{r}_{p-1}|} \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+r_p \end{matrix} \right\}_{\mathbf{r}_p} z^k$ , the identity follows by identification.  $\square$

In [12], we proved the following:

$$\sum_{n \geq 0} B_n(z; \mathbf{r}_p) \frac{t^n}{n!} = B_0(z \exp(t); \mathbf{r}_p) \exp(z(\exp(t) - 1) + r_p t).$$

The following theorem gives more details on the exponential generating function of the  $\mathbf{r}_p$ -Bell polynomials and will be used later.

**Theorem 9.** *We have*

$$\begin{aligned} \sum_{n \geq 0} B_{n+m}(z; \mathbf{r}_p) \frac{t^n}{n!} &= B_m(z \exp(t); \mathbf{r}_p) \exp(z(\exp(t) - 1) + r_p t) \\ &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) \frac{d^{m+k}}{dt^{m+k}} (\exp(z(\exp(t) - 1) + r_p t)). \end{aligned}$$



*Proof.* Use (1) to get

$$\begin{aligned} \sum_{n \geq 0} B_{n+m}(z; \mathbf{r}_p) \frac{t^n}{n!} &= \sum_{n \geq 0} \left( \exp(-z) \sum_{k \geq 0} P_0(k; \mathbf{r}_p) (k + r_p)^{n+m} \frac{z^k}{k!} \right) \frac{t^n}{n!} \\ &= \exp(-z) \sum_{k \geq 0} P_0(k; \mathbf{r}_p) (k + r_p)^m \frac{z^k \exp((k + r_p)t)}{k!} \\ &= B_m(z \exp(t); \mathbf{r}_p) \exp(z(\exp(t) - 1) + r_p t). \end{aligned}$$

For the second part of the theorem, use Theorem 7 to obtain

$$\begin{aligned} \sum_{n \geq 0} B_{n+m}(z; \mathbf{r}_p) \frac{t^n}{n!} &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) \sum_{n \geq 0} B_{n+m+k}(z; r_p) \frac{t^n}{n!} \\ &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) \frac{d^{m+k}}{dt^{m+k}} \left( \sum_{n \geq 0} B_n(z; r_p) \frac{t^n}{n!} \right) \\ &= \sum_{k=0}^{|\mathbf{r}_{p-1}|} a_k(\mathbf{r}_{p-1}) \frac{d^{m+k}}{dt^{m+k}} (\exp(z(\exp(t) - 1) + r_p t)). \end{aligned}$$

□

Using combinatorial arguments, Spivey [13] established the following identity:

$$B_{n+m} = \sum_{k=0}^n \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n}{k} j^{n-k} B_k,$$

where  $B_n$  is the  $n$ -th Bell number, i.e., the number of ways to partition a set of  $n$  elements into non-empty subsets. After that, Belbachir et al. [1] and Gould et al. [7] showed, using different methods, that the polynomial  $B_{n+m}(z) = B_{n+m}(z; \mathbf{0})$  admits a recurrence relation related to the family  $\{z^i B_j(z)\}$  as follows:

$$B_{n+m}(z) = \sum_{k=0}^n \sum_{j=0}^m \left\{ \begin{matrix} m \\ j \end{matrix} \right\} \binom{n}{k} j^{n-k} z^j B_k(z). \tag{4}$$

Recently, Xu [14] gave a recurrence relation on a large family of Stirling numbers and Mihoubi et al. [11] extended the relation (4) to  $r$ -Bell polynomials as follows:

$$B_{n+m,r}(x) = \sum_{k=0}^n \sum_{j=0}^m \left\{ \begin{matrix} m+r \\ j+r \end{matrix} \right\}_r \binom{n}{k} j^{n-k} x^j B_{k,r}(x). \tag{5}$$

Other recurrence relations are given by Mező [10]. The following theorem generalizes the Identities (4) and (5), and the Carlitz's identities [4, 5] given by

$$B_{n+m}(1; r) = \sum_{k=0}^m \left\{ \begin{matrix} m+r \\ k+r \end{matrix} \right\}_r B_n(1; k+r),$$

$$B_n(1; r+s) = \sum_{k=0}^s \left[ \begin{matrix} s+r \\ k+r \end{matrix} \right]_r (-1)^{s-k} B_{n+k}(1; r),$$

and shows that  $B_{n+m}(z; \mathbf{r}_p)$  admits  $r$ -Stirling recurrence coefficients in the families of basis

$$\{z^j B_m(z; \mathbf{r}_p + j\mathbf{e}_p) : 0 \leq j \leq n\},$$

$$\{z^j B_{m+i}(z; r+j) : 0 \leq i \leq |\mathbf{r}_{p-1}|, 0 \leq j \leq n\},$$

where  $B_n(1; r)$  is the number of ways to partition a set of  $n$  elements into non-empty subsets such that the  $r$  first elements are in different subsets.

**Theorem 10.** *We have*

$$B_{n+m}(z; \mathbf{r}_p) = \sum_{j=0}^n \left\{ \begin{matrix} n+r_p \\ j+r_p \end{matrix} \right\}_{r_p} z^j B_m(z; \mathbf{r}_p + j\mathbf{e}_p),$$

$$B_{n+m}(z; \mathbf{r}_p) = \sum_{i=0}^{|\mathbf{r}_{p-1}|} \sum_{j=0}^n \left\{ \begin{matrix} n+r_p \\ j+r_p \end{matrix} \right\}_{r_p} a_i(\mathbf{r}_{p-1}) z^j B_{m+i}(z; r_p + j),$$

$$z^n B_m(z; \mathbf{r}_p + n\mathbf{e}_p) = \sum_{j=0}^n \left[ \begin{matrix} n+r_p \\ j+r_p \end{matrix} \right]_{r_p} (-1)^{n-j} B_{m+j}(z; \mathbf{r}_p).$$

*Proof.* Let  $T_m(z; \mathbf{r}_p) := \sum_{n \geq 0} \left( \sum_{j=0}^n \left\{ \begin{matrix} n+r_p \\ j+r_p \end{matrix} \right\}_{r_p} z^j B_m(z; \mathbf{r}_p + j\mathbf{e}_p) \right) \frac{t^n}{n!}$ . The second identity given in [12, Corollary 12] by

$$\exp(z) B_m(z; \mathbf{r}_p + \mathbf{e}_p) = \frac{d}{dz} (\exp(z) B_m(z; \mathbf{r}_p))$$

can be used to get

$$\exp(z) B_m(z; \mathbf{r}_p + j\mathbf{e}_p) = \frac{d^j}{dz^j} (\exp(z) B_m(z; \mathbf{r}_p)). \tag{6}$$

Identity (6) and the exponential generating function of the  $r$ -Stirling numbers (see [2])

$$\sum_{n \geq j} \left\{ \begin{matrix} n+r_p \\ j+r_p \end{matrix} \right\}_{r_p} \frac{t^n}{n!} = \frac{1}{j!} (\exp(t) - 1)^j \exp(r_p t)$$

prove that

$$\begin{aligned} T_m(z; \mathbf{r}_p) &= \sum_{j \geq 0} B_m(z; \mathbf{r}_p + j\mathbf{e}_p) z^j \frac{1}{j!} (\exp(t) - 1)^j \exp(r_p t) \\ &= \exp(r_p t - z) \sum_{j \geq 0} \frac{d^j}{dz^j} (\exp(z) B_m(z; \mathbf{r}_p)) \frac{(z(\exp(t) - 1))^j}{j!}. \end{aligned}$$

Now, by the Taylor-Maclaurin expansion we have

$$\sum_{j \geq 0} \frac{d^j}{dz^j} (\exp(z) B_m(z; \mathbf{r}_p)) \frac{(u - z)^j}{j!} = \exp(u) B_m(u; \mathbf{r}_p).$$

So, this identity and Theorem 9 show that we have

$$T_m(z; \mathbf{r}_p) = \exp(r_p t - z) \exp(z \exp(t)) B_m(z \exp(t); \mathbf{r}_p) = \sum_{n \geq 0} B_{n+m}(z; \mathbf{r}_p) \frac{t^n}{n!}.$$

By comparing the coefficients of  $t^n$  in the two expressions of  $T_m(z; \mathbf{r}_p)$ , the first identity of this theorem follows. The second identity follows by replacing  $B_m(z; \mathbf{r}_p + j\mathbf{e}_p)$ , as given by its expression in Theorem 7 by

$$\sum_{i=0}^{|\mathbf{r}_{p-1}|} a_i(\mathbf{r}_{p-1}) B_{m+i}(z; j + r_p).$$

For the third identity, let  $A := \sum_{j=0}^n \begin{bmatrix} n+r_p \\ j+r_p \end{bmatrix}_{r_p} (-1)^{n-j} B_{m+j}(z; \mathbf{r}_p)$ . We use Identity (1) and the known identity  $(k + r_p)^{\underline{n}} = \sum_{j=0}^n \begin{bmatrix} n+r_p \\ j+r_p \end{bmatrix}_{r_p} k^j$  (see [2]) to obtain

$$\begin{aligned} A &= \exp(-z) \sum_{k \geq 0} P_m(k; \mathbf{r}_p) \frac{z^k}{k!} \sum_{j=0}^n \begin{bmatrix} n+r_p \\ j+r_p \end{bmatrix}_{r_p} (-1)^{n-j} (k+r_p)^j \\ &= (-1)^n \exp(-z) \sum_{k \geq 0} P_m(k; \mathbf{r}_p) \frac{z^k}{k!} \sum_{j=0}^n \begin{bmatrix} n+r_p \\ j+r_p \end{bmatrix}_{r_p} (-k-r_p)^j \\ &= (-1)^n \exp(-z) \sum_{k \geq 0} P_m(k; \mathbf{r}_p) \frac{z^k}{k!} (-k-r_p+r_p)^{\overline{n}} \\ &= \exp(-z) \sum_{k \geq n} P_m(k; \mathbf{r}_p) k^{\underline{n}} \frac{z^k}{k!} \\ &= z^n \exp(-z) \sum_{k \geq 0} P_m(k+n; \mathbf{r}_p) \frac{z^k}{k!} \\ &= z^n B_m(z; \mathbf{r}_p + n\mathbf{e}_p). \quad \square \end{aligned}$$

As consequences of Theorem 10, some identities for the  $\mathbf{r}_p$ -Stirling numbers of the second kind can be deduced as is shown by the following corollary.

**Corollary 11.** *We have*

$$\begin{aligned} & \sum_{i=0}^k \left\{ \begin{matrix} m + |\mathbf{r}_p| \\ i + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left\{ \begin{matrix} n + r_p \\ k - i + r_p \end{matrix} \right\}_{r_p} = \left\{ \begin{matrix} n + m + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p}, \\ & \sum_{j=0}^n \left\{ \begin{matrix} m + j + |\mathbf{r}_p| \\ k + n + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j} = \left\{ \begin{matrix} m + |\mathbf{r}_p| + n \\ k + r_p + n \end{matrix} \right\}_{\mathbf{r}_p + n\mathbf{e}_p}, \\ & \sum_{j=0}^n \left\{ \begin{matrix} m + j + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j} = 0, \quad k < n. \end{aligned}$$

*Proof.* From the first identity of Theorem 10 we have

$$B_{n+m}(z; \mathbf{r}_p) = \sum_{j=0}^n \left\{ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right\}_{r_p} z^j B_m(z; \mathbf{r}_p + j\mathbf{e}_p)$$

which can be written as

$$\begin{aligned} \sum_{k=0}^{n+m+|\mathbf{r}_p-1|} \left\{ \begin{matrix} n + m + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} z^k &= \sum_{j=0}^n \left\{ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right\}_{r_p} z^j \sum_{i=0}^{m+|\mathbf{r}_p-1|} \left\{ \begin{matrix} m + |\mathbf{r}_p| \\ i + r_p \end{matrix} \right\}_{\mathbf{r}_p} z^i \\ &= \sum_{k=0}^{n+m+|\mathbf{r}_p-1|} z^k \sum_{i=0}^k \left\{ \begin{matrix} m + |\mathbf{r}_p| \\ i + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left\{ \begin{matrix} n + r_p \\ k - i + r_p \end{matrix} \right\}_{r_p}. \end{aligned}$$

Then, the desired identity follows by comparing the coefficients of  $z^k$  in the last expansion. Using the definition of  $B_n(z; \mathbf{r}_p)$  and the third identity of Theorem 10, the second and the third identities of the corollary follow from the definition

$$B_m(z; \mathbf{r}_p + n\mathbf{e}_p) = \sum_{k=0}^{m+|\mathbf{r}_p-1|} \left\{ \begin{matrix} m + |\mathbf{r}_p| + n \\ k + r_p + n \end{matrix} \right\}_{\mathbf{r}_p + n\mathbf{e}_p} z^k$$

and the expansion

$$\begin{aligned} B_m(z; \mathbf{r}_p + n\mathbf{e}_p) &= z^{-n} \sum_{j=0}^n \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j} B_{m+j}(z; \mathbf{r}_p) \\ &= z^{-n} \sum_{j=0}^n \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j} \sum_{k=0}^{m+j+|\mathbf{r}_p-1|} \left\{ \begin{matrix} m + j + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} z^k \\ &= \sum_{k=0}^{m+n+|\mathbf{r}_p-1|} z^{k-n} \sum_{j=0}^n \left\{ \begin{matrix} m + j + |\mathbf{r}_p| \\ k + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j} \\ &= \sum_{k=-n}^{m+|\mathbf{r}_p-1|} z^k \sum_{j=0}^n \left\{ \begin{matrix} m + j + |\mathbf{r}_p| \\ k + n + r_p \end{matrix} \right\}_{\mathbf{r}_p} \left[ \begin{matrix} n + r_p \\ j + r_p \end{matrix} \right]_{r_p} (-1)^{n-j}. \end{aligned}$$

□

### 4. Ordinary Generating Functions

The *ordinary generating function* of the  $r$ -Stirling numbers of the second kind [2] is given by

$$\sum_{n \geq k} \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r t^n = t^k \prod_{j=0}^k (1 - (r+j)t)^{-1}. \tag{7}$$

An analogous result for the  $\mathbf{r}_p$ -Stirling numbers is given by the following theorem.

**Theorem 12.** *Let*

$$\tilde{B}_n(z; \mathbf{r}_p) := \sum_{k=0}^n \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+|\mathbf{r}_p| \end{matrix} \right\}_{\mathbf{r}_p} z^k.$$

*Then, we have*

$$\begin{aligned} \sum_{n \geq k} \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+|\mathbf{r}_p| \end{matrix} \right\}_{\mathbf{r}_p} t^n &= t^{k+|\mathbf{r}_{p-1}|} \left(\frac{1}{t}\right)^{r_1} \cdots \left(\frac{1}{t}\right)^{r_{p-1}} \prod_{j=0}^{k+|\mathbf{r}_{p-1}|} (1 - (r_p + j)t)^{-1}, \\ \sum_{n \geq 0} \tilde{B}_n(z; \mathbf{r}_p) t^n &= \left(\frac{1}{t}\right)^{r_1} \cdots \left(\frac{1}{t}\right)^{r_{p-1}} \sum_{k \geq |\mathbf{r}_{p-1}|} \frac{z^{k-|\mathbf{r}_{p-1}|} t^k}{\prod_{j=0}^k (1 - (r_p + j)t)}. \end{aligned}$$

*Proof.* Use Corollary 8 to obtain

$$\begin{aligned} \sum_{n \geq k} \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+|\mathbf{r}_p| \end{matrix} \right\}_{\mathbf{r}_p} t^n &= \sum_{n \geq k} \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+|\mathbf{r}_{p-1}|+r_p \end{matrix} \right\}_{\mathbf{r}_p} t^n \\ &= \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) t^{-j} \sum_{n \geq k} \left\{ \begin{matrix} n+j+r_p \\ k+|\mathbf{r}_{p-1}|+r_p \end{matrix} \right\}_{\mathbf{r}_p} t^{n+j} \\ &= \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) t^{-j} \sum_{n \geq k+j} \left\{ \begin{matrix} n+r_p \\ k+|\mathbf{r}_{p-1}|+r_p \end{matrix} \right\}_{\mathbf{r}_p} t^n, \end{aligned}$$

and because  $\left\{ \begin{matrix} n+r_p \\ k+|\mathbf{r}_{p-1}|+r_p \end{matrix} \right\}_{\mathbf{r}_p} = 0$  for  $n = k, \dots, k + |\mathbf{r}_{p-1}| - 1$ , we get

$$\sum_{n \geq k} \left\{ \begin{matrix} n+|\mathbf{r}_p| \\ k+|\mathbf{r}_p| \end{matrix} \right\}_{\mathbf{r}_p} t^n = \left( \sum_{n \geq k+|\mathbf{r}_{p-1}|} \left\{ \begin{matrix} n+r_p \\ k+|\mathbf{r}_{p-1}|+r_p \end{matrix} \right\}_{\mathbf{r}_p} t^n \right) \left( \sum_{j=0}^{|\mathbf{r}_{p-1}|} a_j(\mathbf{r}_{p-1}) t^{-j} \right).$$

The first generating function of the theorem follows by using (3) and (7). For the second one, use the definition of  $\tilde{B}_n(z; \mathbf{r}_p)$  and the last expansion.  $\square$

**Acknowledgments** The authors would like to acknowledge the support from the RECITS's laboratory and the PNR project 8/U160/3172. The authors also wish to thank the referee for reading and evaluating the paper thoroughly.

## References

- [1] H. Belbachir, M. Mihoubi, A generalized recurrence for Bell polynomials: An alternate approach to Spivey and Gould Quaintance formulas. *European J. Combin.* **30** (2009), 1254–1256.
- [2] A. Z. Broder, The  $r$ -Stirling numbers. *Discrete Math.* **49** (1984), 241–259.
- [3] J. N. Darroch, On the distribution of the number of successes in independent trials, *Ann. Math. Stat.* (1964), 1317–1321.
- [4] L. Carlitz, Weighted Stirling numbers of the first and second kind – I. *Fibonacci Quart.* **18** (1980), 147–162.
- [5] L. Carlitz, Weighted Stirling numbers of the first and second kind – II. *Fibonacci Quart.* **18** (1980), 242–257.
- [6] G. H. Hardy, J. E. Littlewood, G. Ploya, *Inequalities* (Cambridge: The University Press, 1952).
- [7] H. W. Gould, J. Quaintance, Implications of Spivey's Bell number formula, *J. Integer Seq.* **11** (2008), Article 08.3.7.
- [8] I. Mező, The  $r$ -Bell numbers. *J. Integer Seq.* **14** (2011), Article 11.1.1.
- [9] I. Mező, On the maximum of  $r$ -Stirling numbers. *Adv. Applied Math.* **41** (2008), 293–306.
- [10] I. Mező, The Dual of Spivey's Bell Number Formula, *J. Integer Seq.* **15**(2) (2012), Article 12.2.4.
- [11] M. Mihoubi, H. Belbachir, Linear recurrences for  $r$ -Bell polynomials. Preprint.
- [12] M. Mihoubi, M. S. Maamra, The  $(r_1, \dots, r_p)$ -Stirling numbers of the second kind. *Integers* **12** (2012), Article A35.
- [13] M. Z. Spivey, A generalized recurrence for Bell numbers. *J. Integer Seq.* **11** (2008), Article 08.2.5.
- [14] A. Xu, Extensions of Spivey's Bell number formula. *Electron. J. Comb.* **19** (2) (2012), Article P6.
- [15] F. Z. Zhao, On log-concavity of a class of generalized Stirling numbers. *Electron. J. Comb.* **19** (2) (2012), Article P11.