

4.2 The Mean Value Theorem

Marius Ionescu

11/1/10


Fact

Let f be a function that satisfies the following three hypotheses:


- 1 f is continuous on the closed interval $[a, b]$.
- 2 f is differentiable on the open interval (a, b) .
- 3 $f(a) = f(b)$

Then, there is a number c in (a, b) such that $f'(c) = 0$.

Examples




(a)



(b)

Examples



Example

- Prove that the equation $x^3 + x - 1 = 0$ has exactly one real root.
- Show that the equation $2x - 1 - \sin x = 0$ has exactly one real root.

The Mean Value Theorem

Fact

Let f be a function that fulfills two hypotheses:

- 1 f is continuous on the closed interval $[a, b]$.
- 2 f is differentiable on the open interval (a, b) .


Then, there is a number c in (a, b) such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

or, equivalently,

$$f(b) - f(a) = f'(c)(b - a).$$

Example



Example

Find all numbers c that satisfy the conclusion of the Mean Value Theorem for

- $f(x) = x^3 + x - 1$, $[0, 2]$.
- $f(x) = e^{-2x}$, $[0, 3]$.

Example

Example

Suppose that $f(0) = -3$ and $f'(x) \leq 5$ for all values of x . How large can $f(2)$ possibly be?