

2.2 The Limit of a Function

Marius Ionescu

09/02/2010

The Legacy of Galileo, Newton, and Leibniz

- Galileo
 - was interested in falling bodies.
 - forged a new scientific methodology - *observe nature, construct experiments to test what you observe, and construct theories that explain the observations.*

The Legacy of Galileo, Newton, and Leibniz

- Galileo
 - was interested in falling bodies.
 - forged a new scientific methodology - *observe nature, construct experiments to test what you observe, and construct theories that explain the observations.*
- Newton
 - was able, using his new tools of calculus, to explain why falling bodies behave in this way: an object, falling under the influence of gravity, will have constant acceleration of $9.8m/sec^2$.
 - his laws of motion and of universal gravitation drew under one simple mathematical theory Newton's laws of falling bodies, Kepler's laws of planetary motion, the motion of a simple pendulum, and virtually every other instance of dynamic motion observed in the universe.

The Legacy of Galileo, Newton, and Leibniz

- Galileo
 - was interested in falling bodies.
 - forged a new scientific methodology - *observe nature, construct experiments to test what you observe, and construct theories that explain the observations.*
- Newton
 - was able, using his new tools of calculus, to explain why falling bodies behave in this way: an object, falling under the influence of gravity, will have constant acceleration of $9.8m/sec^2$.
 - his laws of motion and of universal gravitation drew under one simple mathematical theory Newton's laws of falling bodies, Kepler's laws of planetary motion, the motion of a simple pendulum, and virtually every other instance of dynamic motion observed in the universe.
- Leibnitz
 - co-inventor of calculus, took a slightly different point of view but also studied rates of change in a general setting.


- How do we find the velocity of a moving object at time t ?


- How do we find the velocity of a moving object at time t ?
- What in fact do we mean by *velocity* of the object at the instant of time t ?? We know how to find the average velocity of an object during a time interval $[t_1, t_2]$?

Definition

We say that a function f approaches the limit L as x approaches a , written $\lim_{x \rightarrow a} f(x) = L$, if we can make $f(x)$ as close to L as we please by taking x sufficiently close to a .

Example





Definition

We write

$$\lim_{x \rightarrow a^-} f(x) = L$$

and say the left-hand limit of $f(x)$ as x approaches a –or the limit of $f(x)$ as x approaches a from the left– is equal to L if we can make the values of $f(x)$ arbitrarily close to L by taking x to be sufficiently close to a and x less than a .


Fact

The limit of f as $x \rightarrow a$ exists if and only if both the right-hand and left-hand limits exist and have the same value. i.e.

$$\lim_{x \rightarrow a} f(x) = L \Leftrightarrow \lim_{x \rightarrow a^-} f(x) = L \text{ and } \lim_{x \rightarrow a^+} f(x) = L.$$

Infinite Limits

Compute the limit $\lim_{x \rightarrow 0} 1/x^2$.



Definition


Let f be a function defined on both sides of a , except possibly at a itself. Then,

$$\lim_{x \rightarrow a} f(x) = \infty$$

means that the values of $f(x)$ can be made arbitrarily large—as large as we please—by taking x sufficiently close to a , but not equal to a .

Example

Evaluate $\lim_{x \rightarrow \pi/2} \tan x$.



Definition

The line $x = a$ is called a **vertical asymptote** of the curve $y = f(x)$ if at least one of the following statement is true:

$$\begin{array}{lll} \lim_{x \rightarrow a} f(x) = \infty & \lim_{x \rightarrow a^+} f(x) = \infty & \lim_{x \rightarrow a^-} f(x) = \infty \\ \lim_{x \rightarrow a} f(x) = -\infty & \lim_{x \rightarrow a^+} f(x) = -\infty & \lim_{x \rightarrow a^-} f(x) = -\infty \end{array}$$