


## 2.7 Derivatives and Rates of Change

Marius Ionescu

September 15, 2010


# The Tangent Line and Their Slope

- **The Tangent Line Problem** Given a function  $y = f(x)$  defined in an open interval and a point  $x_0$  in the interval, define the tangent line at the point  $(x_0, f(x_0))$  on the graph of  $f$ .




# Example

Find the equations of the tangent lines to the graph of  $f(x) = \sqrt{1-x^2}$  at the points  $(0, 1)$  and  $(\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})$ .



# Example

Let  $f(x) = x^2$ .



# The slope of the tangent

## Definition

Given a function  $f$  and a point  $x_0$  in its domain, the slope of the tangent line at the point  $(x_0, f(x_0))$  on the graph of  $f$  is


$$m = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}.$$

## Example

Given  $f(x) = \sqrt{x}$ , find the equation of the tangent line at  $x = 4$ .

# Example

Find the tangent line to the graph of  $f(x) = x^{1/3}$  at  $x = 0$ .




# Example

Let  $f$  be the piecewise defined function

$$f(x) = \begin{cases} 2 - x^2 & x \leq 1 \\ x^3 & x > 1 \end{cases}$$

Is the function continuous, and does it have a tangent line at  $x = 1$ ?





## Definition

The **derivative of a function  $f$  at a number  $a$** , denoted by  $f'(a)$ , is

$$f'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}.$$

Equivalently

$$f'(a) = \lim_{x \rightarrow a} \frac{f(x) - f(a)}{x - a}.$$

# The tangent line (revisited)

## Fact

*Tangent line* The tangent line to  $y = f(x)$  at  $(a, f(a))$  is the line through  $(a, f(a))$  whose slope is equal to  $f'(a)$ , the derivative of  $f$  at  $a$ .

## Definition

- The **average rate of change of  $y$  with respect to  $x$**  over the interval  $[x_1, x_2]$  is

$$\frac{\Delta y}{\Delta x} = \frac{f(x_2) - f(x_1)}{x_2 - x_1}.$$

- The **instantaneous rate of change of  $y$  with respect to  $x$**  is

$$\lim_{\Delta x \rightarrow 0} \frac{\Delta y}{\Delta x} = \lim_{x_2 \rightarrow x_1} \frac{f(x_2) - f(x_1)}{x_2 - x_1}.$$

## Example

A manufacturer produces bolts of a fabric with a fixed width. The cost of producing  $x$  yards of this fabric is  $C = f(x)$  dollars.

- 1 What is the meaning of the derivative  $f'(x)$ ? What are its units?
- 2 In practical terms, what does it mean to say that  $f'(1,000) = 9$ ?
- 3 Which do you think is greater,  $f'(50)$  or  $f'(500)$ ? What about  $f'(5,000)$ ?