

# Sections 13.1 and 13.2: Vector Functions, Space Curves, and Derivatives

Marius Ionescu

September 21, 2012

# Vector Functions

- A vector function is a function  $\mathbf{r}(t)$  whose domain is the set of real numbers and whose range is a set of vectors – in general three-dimensional vectors.
- If  $f(t)$ ,  $g(t)$ , and  $h(t)$  are the components of the vector  $\mathbf{r}(t)$ , then they are called the **component functions** of  $\mathbf{r}$ .
- We write

$$\begin{aligned}\mathbf{r}(t) &= \langle f(t), g(t), h(t) \rangle \\ &= f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}\end{aligned}$$

## Limit of a vector function

### Fact

If  $\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle$ , then

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \rangle$$

## Example

### Example


Find the limit

$$\lim_{t \rightarrow \pi/4} \langle \cos t, \sin t, t \rangle$$

# Space Curves

## Example


$$\mathbf{r}(t) = \langle t^2, \ln(4-t), \sqrt{t} \rangle$$



# The Helix

## Example


$$r(t) = \cos t \mathbf{i} + \sin t \mathbf{j} + t \mathbf{k}$$



# The Toroidal

## Example


$$\mathbf{r}(t) = (4 + \sin(2t)) \cos t \mathbf{i} + (4 + \sin 20t) \sin t \mathbf{j} + \cos 20t \mathbf{k}$$



## Example

### Example

Find a vector function that represents the curve of intersection of the cylinder  $x^2 + y^2 = 1$  and the plane  $y + z = 1$ .





# Derivatives of Vector Functions

- The derivative  $\mathbf{r}'$  of  $\mathbf{r}$  is

$$\frac{d\mathbf{r}}{dt} = \mathbf{r}'(t) = \lim_{h \rightarrow 0} \frac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$$

if the limit exists.

- The vector  $\mathbf{r}'(t)$  is called the **tangent vector**.
- The **unit tangent vector** is

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$$

## How to compute $\mathbf{r}'(t)$ ?

- If  $\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle$ , where  $f$ ,  $g$ , and  $h$  are differentiable functions, then

$$\mathbf{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle$$

## Examples

### Examples

- Find the derivative of  $\mathbf{r}(t) = (2t + t^2)\mathbf{i} + e^{-t^2}\mathbf{j} + \sin(2t)\mathbf{k}$ .
- Find parametric equations for the tangent line to the helix with parametric equations

$$x = 2 \cos t, \quad y = \sin t, \quad z = t$$

at the point  $(0, 1, \pi/2)$ .

# Differentiation Rules

$$[\mathbf{u}(t) + \mathbf{v}(t)]' = \mathbf{u}'(t) + \mathbf{v}'(t)$$

$$(c\mathbf{u}(t))' = c\mathbf{u}'(t)$$

$$(f(t)\mathbf{u}(t))' = f'(t)\mathbf{u}(t) + f(t)\mathbf{u}'(t)$$

$$(\mathbf{u}(t) \cdot \mathbf{v}(t))' = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$(\mathbf{u}(t) \times \mathbf{v}(t))' = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

$$(\mathbf{u}(f(t)))' = f'(t)\mathbf{u}'(f(t)).$$