

## 14.3: Partial Derivatives

Marius Ionescu

October 12, 2012

# Partial derivative of $f$ with respect to $x$

Definition

# Partial derivative of $f$ with respect to $x$

## Definition

- Let  $f(x, y)$  be a function of two variables.

# Partial derivative of $f$ with respect to $x$

## Definition

- Let  $f(x, y)$  be a function of two variables.
- Let  $y = b$  be fixed.

# Partial derivative of $f$ with respect to $x$

## Definition

- Let  $f(x, y)$  be a function of two variables.
- Let  $y = b$  be fixed.
- Then  $g(x) = f(x, b)$  is a function of a single variable  $x$ .

## Partial derivative of $f$ with respect to $x$

### Definition

- Let  $f(x, y)$  be a function of two variables.
- Let  $y = b$  be fixed.
- Then  $g(x) = f(x, b)$  is a function of a single variable  $x$ .
- If  $g$  has a derivative at  $a$ , then we call it the **partial derivative of  $f$  with respect to  $x$  at  $(a, b)$**

$$f_x(a, b) = g'(a)$$

# Partial derivative of $f$ with respect to $y$

Definition

# Partial derivative of $f$ with respect to $y$

## Definition

- Now keep  $x = a$  fix.



# Partial derivative of $f$ with respect to $y$

## Definition

- Now keep  $x = a$  fix.
- Let  $h(y) = f(a, y)$ .

# Partial derivative of $f$ with respect to $y$

## Definition

- Now keep  $x = a$  fix.
- Let  $h(y) = f(a, y)$ .
- If  $h$  has a derivative at  $b$ , then we call it the **partial derivative of  $f$  with respect to  $y$  at  $(a, b)$**

$$f_y(a, b) = h'(b)$$

# Partial Derivatives

# Partial Derivatives

- By the definition of a derivative, we have

$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a + h, b) - f(a, b)}{h}$$

$$f_y(a, b) = \lim_{h \rightarrow 0} \frac{f(a, b + h) - f(a, b)}{h}$$

# Partial Derivatives

- By the definition of a derivative, we have

$$f_x(a, b) = \lim_{h \rightarrow 0} \frac{f(a + h, b) - f(a, b)}{h}$$

$$f_y(a, b) = \lim_{h \rightarrow 0} \frac{f(a, b + h) - f(a, b)}{h}$$

- The partial derivatives of  $f(x, y)$  are the functions  $f_x(x, y)$  and  $f_y(x, y)$  obtained by letting the point  $(a, b)$  vary.

# Notations

- If  $z = f(x, y)$ , we write

$$f_x(x, y) = f_x = \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} f(x, y) = \frac{\partial z}{\partial x} = f_1 = D_1 f = D_x f$$
$$f_y(x, y) = f_y = \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} f(x, y) = \frac{\partial z}{\partial y} = f_2 = D_2 f = D_y f$$

# Rule for Finding Partial Derivatives of $z = f(x, y)$

## Rule for Finding Partial Derivatives of $z = f(x, y)$

- To find  $f_x$  regard  $y$  as a constant and differentiate  $f(x, y)$  with respect to  $x$ .



## Rule for Finding Partial Derivatives of $z = f(x, y)$

- To find  $f_x$  regard  $y$  as a constant and differentiate  $f(x, y)$  with respect to  $x$ .
- To find  $f_y$  regard  $x$  as a constant and differentiate  $f(x, y)$  with respect to  $y$ .

# Examples

## Examples

# Examples

## Examples

- If  $f(x, y) = x^2 + 3x^3y - xy^2$  find  $f_x(0, 1)$  and  $f_y(1, 0)$

# Examples

## Examples

- If  $f(x, y) = x^2 + 3x^3y - xy^2$  find  $f_x(0, 1)$  and  $f_y(1, 0)$
- Find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  for the functions

# Examples

## Examples

- If  $f(x, y) = x^2 + 3x^3y - xy^2$  find  $f_x(0, 1)$  and  $f_y(1, 0)$
- Find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  for the functions

•

$$f(x, y) = \frac{2y}{y + \cos x}$$

# Examples

## Examples

- If  $f(x, y) = x^2 + 3x^3y - xy^2$  find  $f_x(0, 1)$  and  $f_y(1, 0)$
- Find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  for the functions

- 

$$f(x, y) = \frac{2y}{y + \cos x}$$

- 

$$f(x, y) = e^{x^2+y^2+1}$$

# Examples

## Examples

- If  $f(x, y) = x^2 + 3x^3y - xy^2$  find  $f_x(0, 1)$  and  $f_y(1, 0)$
- Find  $\frac{\partial f}{\partial x}$  and  $\frac{\partial f}{\partial y}$  for the functions

- 

$$f(x, y) = \frac{2y}{y + \cos x}$$

- 

$$f(x, y) = e^{x^2+y^2+1}$$

- 

$$f(x, y) = \ln(x + y)$$

## Examples (cont'd)

### Example

- Find  $\frac{\partial z}{\partial x}$  and  $\frac{\partial z}{\partial y}$  if  $z$  is defined implicitly as a function of  $x$  and  $y$  by the equation

$$x^3 + y^3 + z^3 + 6xyz = 1.$$



# Interpretations of Partial Derivatives

# Interpretations of Partial Derivatives

- Partial derivative can be interpreted as rates of change.

# Interpretations of Partial Derivatives

- Partial derivative can be interpreted as rates of change.
- The geometric interpretation: the partial derivatives are the slopes of the tangent lines at  $P(a, b, c)$  to the curves given by the intersection of the surface given by  $z = f(x, y)$  and the planes  $x = a$  and  $y = b$ .

# Higher Derivatives

## Definition

# Higher Derivatives

## Definition

- If  $f$  is a function of two variables, then its partial derivatives  $f_x$  and  $f_y$  are also functions of two variables.

# Higher Derivatives

## Definition

- If  $f$  is a function of two variables, then its partial derivatives  $f_x$  and  $f_y$  are also functions of two variables.
- So why stop here?

# Higher Derivatives

## Definition

- If  $f$  is a function of two variables, then its partial derivatives  $f_x$  and  $f_y$  are also functions of two variables.
- So why stop here?
- The **second partial derivatives** of  $f$  are

$$(f_x)_x = f_{xx} = f_{11} = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial x} \right) = \frac{\partial^2 f}{\partial^2 x} = \frac{\partial^2 z}{\partial^2 x}$$

$$f_{xy} = \frac{\partial^2 f}{\partial x \partial y} = \dots$$

$$f_{yx} = \frac{\partial^2 f}{\partial y \partial x} = \dots$$

$$f_{yy} = \frac{\partial^2 f}{\partial^2 y}$$

## Example

### Example

- Find the second derivatives of

$$f(x, y) = x^3 + x^2y^3 - 2y^2$$



# Clairaut's Theorem

## Theorem

- Suppose  $f$  is defined on a disk  $D$  that contains the point  $(a, b)$ . If the functions  $f_{xy}$  and  $f_{yx}$  are both continuous on  $D$ , then

$$f_{xy}(a, b) = f_{yx}(a, b)$$

# Partial derivatives of order 3 and higher

## Examples

# Partial derivatives of order 3 and higher

## Examples

- Calculate  $f_{xxy}$  if  $f(x, y) = \sin(3x^2 + xy)$ .

# Partial derivatives of order 3 and higher

## Examples

- Calculate  $f_{xxy}$  if  $f(x, y) = \sin(3x^2 + xy)$ .
- Find the partial derivatives of

$$f(x, y) = \int_x^y e^{t^2+t+1} dt$$

# Partial derivatives of order 3 and higher

## Examples

- Calculate  $f_{xxy}$  if  $f(x, y) = \sin(3x^2 + xy)$ .
- Find the partial derivatives of

$$f(x, y) = \int_x^y e^{t^2+t+1} dt$$

- Find  $f_x, f_y, f_{xy}, f_{yx}$  for

$$f(x, y) = xye^{3xy}$$