

Equations of the tangent plane

Notes

Definition

- Suppose f has a continuous partial derivatives.
- An equation of the tangent plane to the surface z = f(x, y) at the point $P(x_0, y_0, z_0)$ is

$$z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

Marius Ionescu () 14.4: Tangent Planes and Linear Approxi

October 15, 2012 3 / 13

Examples

Example

• Find the tangent plane to the elliptic paraboloid $z = 2x^2 + y^2$ at the point (1, 1, 3).



Notes







Linear Approximations

Definition

• The linear function whose graph is this tangent plane

 $L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$

is called the **linearization** of f at (a, b) and the approximation

$$f(x,y) \approx f(a,b) + f_x(a,b)(x-a) + f_y(a,b)(y-b)$$

is called the linear approximation or the tangent plane approximation of f at (a, b).

Notes

Notes

5 / 13

Examples

Notes

Examples

- Find the linearization of the function $f(x, y) = \sqrt{xy}$ at the point (4, 16).
- Find the linearization of the function $f(x, y) = 1 + y + x \cos y$ at $P_0(0, 0)$.

Marius Ionescu () 14.4: Tangent Planes and Linear Approxi

The increment of z

Definition

• Recall that for a function of one variable, y = f(x), if x changes from a to $a + \Delta x$, we defined the increment of y as

$$\Delta y = f(a + \Delta x) - f(a).$$

• If f is differentiable at a, then

$$\Delta y = f'(a)\Delta x + \epsilon \Delta x$$

where $\epsilon \rightarrow 0$ as $\Delta x \rightarrow 0$.

Notes

October 15, 2012 7 / 13

The increment of z

Notes

Notes

Definition

• If z = f(x, y) and x changes from (a, b) to $(a + \Delta x, b + \Delta y)$, then the **increment** of z is

$$\Delta z = f(a + \Delta x, b + \Delta y) - f(a, b)$$

• If z = f(x, y), then f is differentiable at (a, b) if Δz can be expressed in the form

$$\Delta z = f_x(a, b)\Delta x + f_y(a, b)\Delta y + \epsilon_1 \Delta x + \epsilon_2 \Delta y,$$

where ϵ_1 and $\epsilon_2 \rightarrow 0$ as $(\Delta x, \Delta y) \rightarrow (0, 0)$.

Marius Ionescu ()	14.4: Tangent Planes and Linear Approxi	October 15, 2012	9 / 13
Fact			1
		\ <i>\</i>	- 1
If the partial der	vatives f_x and f_y exist near (a, b)) and are contini	ious
at (a, b), then f	is differentiable at (a, b).		- 1

Example

• Show that $f(x, y) = xe^{xy}$ is differentiable at (1,0) and find its linearization there. Then use it to approximate f(1.1, -0.1).

Marius Ionescu () 14.4: Tangent Planes and Linear Approxi October 15, 2012 11 / 13

Differentials

Definition

• For a differentiable function z = f(x, y) we define the differential dz, also called the total differential, is defined by

$$dz = f_x(x, y)dx + f_y(x, y)dy = \frac{\partial z}{\partial x}dx + \frac{\partial z}{\partial y}dy,$$

where the differentials dx and dy are independent variables.

• If $dx = \Delta x = x - a$ and $dy = \Delta y = y - b$ the the differential of z is

$$dz = f_x(a,b)(x-a) + f_y(a,b)(y-b)$$

Notes

Examples

Examples

- If $f(x, y) = x^2 + 3xy y^2$, find the differential dz.
- If x changes from 2 to 2.05 and y changes 3 to 2.96, compare the values of Δz and dz.

Marius Ionescu () 14.4: Tangent Planes and Linear Approxi

October 15, 2012 13 / 13

Notes