



**LINEAR RECURRENCES FROM COUNTING SCHREIER-TYPE  
MULTISETS**

**Hùng Việt Chu**

*Department of Mathematics, Washington and Lee University, Lexington, Virginia*  
hchu@wlu.edu

**Yubo Geng**

*College of Liberal Arts, University of Minnesota, Minneapolis, Minnesota*  
geng0114@umn.edu

**Julian King**

*Department of Mathematics, SUNY Geneseo, Geneseo, New York*  
jkh12@geneseo.edu

**Steven J. Miller**

*Dept. of Math. and Stats., Williams College, Williamstown, Massachusetts*  
sjm1@williams.edu

**Garrett Tresch**

*Department of Mathematics, Texas A&M University, College Station, Texas*  
treschgd@tamu.edu

**Zachary Louis Vasseur**

*Department of Mathematics, Texas A&M University, College Station, Texas*  
zachary.l.v@tamu.edu

*Received: 9/10/25, Accepted: 3/24/26, Published: 5/1/26*

**Abstract**

A nonempty set  $F$  is Schreier if  $\min F \geq |F|$ . Bird observed that counting Schreier sets in a certain way produces the Fibonacci sequence. Since then, various connections between variants of Schreier sets and well-known sequences have been discovered. Building on these works, we prove a linear recurrence for the sequence that counts multisets  $F$  with  $\min F \geq p|F|$ . In particular, if we let

$$\mathcal{A}_{p,n}^{(s)} := \{F \subset \underbrace{\{1, \dots, 1\}}_s, \dots, \underbrace{\{n-1, \dots, n-1\}}_s, n\} : n \in F \text{ and } \min F \geq p|F|\},$$

then  $|\mathcal{A}_{p,n}^{(s)}| = \sum_{i=0}^s |\mathcal{A}_{p,n-1-ip}^{(s)}|$ . If we color  $s$  copies of the same integer by different colors from 1 to  $s$ , i.e.,  $\mathcal{B}_{p,n}^{(s)} := \{F \subset \{1_1, \dots, 1_s, \dots, (n-1)_1, \dots, (n-1)_s, n\} : n \in F \text{ and } \min F \geq p|F|\}$ , then  $|\mathcal{B}_{p,n}^{(s)}| = \sum_{i=0}^s \binom{s}{i} |\mathcal{B}_{p,n-1-ip}^{(s)}|$ . Lastly, we count Schreier sets that do not admit multiples of a given integer  $u \geq 2$  and witness linear recurrences whose coefficients are drawn from the  $u^{\text{th}}$  row of Pascal's triangle and have alternating signs, except possibly the last one.

**1. Introduction**

A finite, nonempty set  $F \subset \mathbb{N}$  is *Schreier* if  $\min F \geq |F|$ . Bird [5] discovered a fascinating connection between Schreier sets and Fibonacci numbers: for  $n \in \mathbb{N}$ ,

$$|\{F \subset \{1, 2, \dots, n\} : F \text{ is Schreier and } n \in F\}| = F_n,$$

where  $(F_n)_{n=1}^\infty$  is the Fibonacci sequence with  $F_1 = F_2 = 1$  and  $F_n = F_{n-1} + F_{n-2}$  for  $n \geq 3$ . If instead, we consider Schreier multisets, [7, Theorem 1] states that

$$|\{F \subset \underbrace{\{1, \dots, 1\}}_s, \dots, \underbrace{\{n-1, \dots, n-1\}}_s, n\} : F \text{ is Schreier and } n \in F\}| = F_n^{(s+1)},$$

where for each  $m \geq 2$ , the sequence  $(F_n^{(m)})_{n=1}^\infty$  is the  $m$ -step Fibonacci sequence with

$$\begin{aligned} F_{2-m}^{(m)} &= \dots = F_0^{(m)} = 0, & F_1^{(m)} &= 1, & \text{and} \\ F_n^{(m)} &= F_{n-1}^{(m)} + F_{n-2}^{(m)} + \dots + F_{n-m}^{(m)}, & \text{for } n &\geq 2. \end{aligned}$$

These  $m$ -step Fibonacci sequences are listed in The On-Line Encyclopedia of Integer Sequences [11] as [A000045](#), [A000073](#), [A000078](#), and so on. For more connections between Schreier-type sets and various sequences, the readers may refer to [1, 2, 3, 8, 9].

Inspired by [2, 7], we investigate the following general problem of counting Schreier-type multisets.

**Problem 1.** Given  $(p, q) \in \mathbb{N}^2$  and for each  $n \in \mathbb{N}$ , a sequence of nonnegative integers  $(s_{n,i})_{i=1}^n$ , define

$$H_{p,q,n}^{(s_{n,i})} := \{F \subset \underbrace{\{1, \dots, 1\}}_{s_{n,1}}, \dots, \underbrace{\{n, \dots, n\}}_{s_{n,n}}\} : n \in F \text{ and } q \min F \geq p|F|\}.$$

Find the recurrence (if any) of  $(|H_{p,q,n}^{(s_{n,i})}|)_{n=1}^\infty$ .

We solve Problem 1 and its colored version when  $q = 1$ ,  $s_{n,1} = \dots = s_{n,n-1} = s$  for some fixed positive integer  $s$ , and  $s_{n,n} = 1$ . Our main results show that the sequences from counting these multisets both satisfy linear recurrences whose indices are in arithmetic progression as well as linear recurrences with coefficients taken from Pascal’s triangle, respectively. For  $(s, p, n) \in \mathbb{N}^3$ , define

$$\mathcal{A}_{p,n}^{(s)} := \{F \subset \underbrace{\{1, \dots, 1\}}_s, \dots, \underbrace{\{n-1, \dots, n-1\}}_s, n\} : n \in F \text{ and } \min F \geq p|F|\}.$$

Table 1 gives sample data for  $(|\mathcal{A}_{p,n}^{(s)}|)_{n=1}^\infty$ .

| $n$                         | 1 | 2 | 3 | 4 | 5 | 6  | 7  | 8  | 9   | 10  | 11  | 12  | OEIS                    |
|-----------------------------|---|---|---|---|---|----|----|----|-----|-----|-----|-----|-------------------------|
| $ \mathcal{A}_{1,n}^{(1)} $ | 1 | 1 | 2 | 3 | 5 | 8  | 13 | 21 | 34  | 55  | 89  | 144 | <a href="#">A000045</a> |
| $ \mathcal{A}_{2,n}^{(1)} $ | 0 | 1 | 1 | 1 | 2 | 3  | 4  | 6  | 9   | 13  | 19  | 28  | <a href="#">A078012</a> |
| $ \mathcal{A}_{1,n}^{(2)} $ | 1 | 1 | 2 | 4 | 7 | 13 | 24 | 44 | 81  | 149 | 274 | 504 | <a href="#">A000073</a> |
| $ \mathcal{A}_{2,n}^{(2)} $ | 0 | 1 | 1 | 1 | 2 | 3  | 5  | 8  | 12  | 19  | 30  | 47  | <a href="#">A060961</a> |
| $ \mathcal{A}_{1,n}^{(3)} $ | 1 | 1 | 2 | 4 | 8 | 15 | 29 | 56 | 108 | 208 | 401 | 773 | <a href="#">A000078</a> |
| $ \mathcal{A}_{2,n}^{(3)} $ | 0 | 1 | 1 | 1 | 2 | 3  | 5  | 8  | 13  | 21  | 33  | 53  | <a href="#">A117760</a> |

Table 1: The first 12 values of  $(|\mathcal{A}_{p,n}^{(s)}|)_{n=1}^\infty$  with different  $(s, p)$ .

**Theorem 1.** For  $1 \leq n \leq sp + 1$ ,

$$|\mathcal{A}_{p,n}^{(s)}| = \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \binom{n - pk + k - 2}{k - 1}.$$

For  $n \geq sp + 2$ , we have

$$|\mathcal{A}_{p,n}^{(s)}| = \sum_{i=0}^s |\mathcal{A}_{p,n-1-ip}^{(s)}|. \tag{1}$$

**Remark 1.** It is interesting to see that  $p$  and  $s$  have different effects on Recurrence (1). While  $(s + 1)$  gives the number of terms in the recurrence,  $p$  equals the gap between consecutive indices.

**Remark 2.** The problem of extending Theorem 1 to all  $q \in \mathbb{N}$  (instead of  $q = 1$ ) remains open and is discussed in the last section.

Next, we give a colored version of Theorem 1. Given an integer  $k$ , we color the  $s$  copies of  $k$  using  $s$  different colors and denote them by  $k_1, k_2, \dots, k_s$ . For example, while  $k_1$  and  $k_2$  have the same numerical value, the two sets  $\{k_1\}$  and  $\{k_2\}$  are distinguishable. Let

$$\mathcal{B}_{p,n}^{(s)} := \{F \subset \{1_1, \dots, 1_s, \dots, (n-1)_1, \dots, (n-1)_s, n\} : n \in F \text{ and } \min F \geq p|F|\}.$$

We collect sample data for  $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^\infty$  in Table 2.

| $n$                         | 1 | 2 | 3 | 4  | 5  | 6  | 7   | 8   | 9    | 10   | 11   | OEIS                    |
|-----------------------------|---|---|---|----|----|----|-----|-----|------|------|------|-------------------------|
| $ \mathcal{B}_{1,n}^{(1)} $ | 1 | 1 | 2 | 3  | 5  | 8  | 13  | 21  | 34   | 55   | 89   | <a href="#">A000045</a> |
| $ \mathcal{B}_{2,n}^{(1)} $ | 0 | 1 | 1 | 1  | 2  | 3  | 4   | 6   | 9    | 13   | 19   | <a href="#">A078012</a> |
| $ \mathcal{B}_{1,n}^{(2)} $ | 1 | 1 | 3 | 6  | 13 | 28 | 60  | 129 | 277  | 595  | 1278 | <a href="#">A002478</a> |
| $ \mathcal{B}_{2,n}^{(2)} $ | 0 | 1 | 1 | 1  | 3  | 5  | 8   | 15  | 26   | 45   | 80   | <a href="#">A193147</a> |
| $ \mathcal{B}_{1,n}^{(3)} $ | 1 | 1 | 4 | 10 | 26 | 69 | 181 | 476 | 1252 | 3292 | 8657 | <a href="#">A099234</a> |
| $ \mathcal{B}_{2,n}^{(3)} $ | 0 | 1 | 1 | 1  | 4  | 7  | 13  | 28  | 53   | 105  | 211  | <a href="#">A373718</a> |

Table 2: The first 11 values of  $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^\infty$  with different  $(s, p)$ .

**Theorem 2.** *Let  $s$  and  $p$  be positive integers. For  $1 \leq n \leq sp + 1$ ,*

$$|\mathcal{B}_{p,n}^{(s)}| = \sum_{k=1}^{\lfloor \frac{ns+1}{ps+1} \rfloor} \binom{(n-kp)s}{k-1}.$$

For  $n \geq sp + 2$ , we have

$$\begin{aligned} |\mathcal{B}_{p,n}^{(s)}| &= \binom{s}{0} |\mathcal{B}_{p,n-1}^{(s)}| + \binom{s}{1} |\mathcal{B}_{p,n-1-p}^{(s)}| + \dots + \binom{s}{s} |\mathcal{B}_{p,n-1-sp}^{(s)}| \\ &= \sum_{i=0}^s \binom{s}{i} |\mathcal{B}_{p,n-1-ip}^{(s)}|. \end{aligned}$$

Motivated by a recent result in [9] that counts Schreier sets of multiples of a fixed  $u \geq 2$ , our last result counts Schreier sets that do not admit any multiple of  $u$ . If  $g_u(n)$  denotes the  $n^{\text{th}}$  positive integer not divisible by  $u$ , then as we show later (Lemma 2),  $g_u(n) = \lfloor (un - 1)/(u - 1) \rfloor$ . Let  $G_{u,n}$  be the set of the first  $n$  positive integers that are not divisible by  $u$  and define

$$\mathcal{D}_{u,n} := \left\{ F \subset G_{u,n} : \left\lfloor \frac{un-1}{u-1} \right\rfloor \in F \text{ and } F \text{ is Schreier} \right\}.$$

| $n$                   | 1 | 2 | 3 | 4 | 5 | 6  | 7  | 8  | 9  | 10  | 11  | 12  | OEIS                    |
|-----------------------|---|---|---|---|---|----|----|----|----|-----|-----|-----|-------------------------|
| $ \mathcal{D}_{2,n} $ | 1 | 1 | 2 | 4 | 7 | 12 | 21 | 37 | 65 | 114 | 200 | 351 | <a href="#">A005251</a> |
| $ \mathcal{D}_{3,n} $ | 1 | 1 | 2 | 3 | 5 | 9  | 16 | 28 | 48 | 81  | 136 | 229 | <a href="#">A137402</a> |
| $ \mathcal{D}_{4,n} $ | 1 | 1 | 2 | 3 | 5 | 8  | 13 | 22 | 38 | 66  | 114 | 195 | N/A                     |
| $ \mathcal{D}_{5,n} $ | 1 | 1 | 2 | 3 | 5 | 8  | 13 | 21 | 34 | 56  | 94  | 160 | N/A                     |
| $ \mathcal{D}_{6,n} $ | 1 | 1 | 2 | 3 | 5 | 8  | 13 | 21 | 34 | 55  | 89  | 145 | N/A                     |

Table 3: The first 12 values of  $(|\mathcal{D}_{u,n}|)_{n=1}^\infty$  with  $u \in \{2, \dots, 6\}$ .

We establish the following recurrence for the sequence  $(|\mathcal{D}_{u,n}|)_{n=1}^\infty$ , which is supported by the data presented in Table 3.

**Theorem 3.** For  $1 \leq n \leq 2u - 1$ , we have  $|\mathcal{D}_{u,n}| = F_n$ . For  $n \geq 2u$ ,

$$|\mathcal{D}_{u,n}| = \sum_{i=1}^u (-1)^{i-1} \binom{u}{i} |\mathcal{D}_{u,n-i}| + |\mathcal{D}_{u,n-2u+1}|.$$

We devote Sections 2, 3, and 4 to the proof of Theorems 1, 2, and 3, respectively. Section 5 discusses directions for future investigations. To keep our proofs concise and emphasize the main ideas, we relegate certain technical proofs to the Appendix.

### 2. Uncolored Generalized Schreier Multisets

We give a formula to compute  $|\mathcal{A}_{p,n}^{(s)}|$ , which is used to prove the first statement of Theorem 1, while we prove the second statement using bijective maps. For  $k \geq 1$ , each  $k$ -element multiset in  $\mathcal{A}_{p,n}^{(s)}$  is uniquely determined by  $k - 1$  numbers from the multiset

$$\underbrace{\{pk, \dots, pk\}}_s, \dots, \underbrace{\{n-1, \dots, n-1\}}_s.$$

Let  $\binom{n}{m}_s$  be the number of ways to distribute  $m$  identical objects among  $n$  labeled boxes, each of which contains at most  $s - 1$  objects. According to [6, (1.16)],

$$\binom{n}{m}_s = \sum_{k=0}^{\lfloor m/s \rfloor} (-1)^k \binom{n}{k} \binom{n+m-sk-1}{n-1}.$$

It follows that the number of  $k$ -element multisets in  $\mathcal{A}_{p,n}^{(s)}$  is  $\binom{n-pk}{k-1}_{s+1}$  under the condition  $s(n - pk) \geq k - 1$ . Therefore,

$$\begin{aligned} |\mathcal{A}_{p,n}^{(s)}| &= \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \binom{n-pk}{k-1}_{s+1} \\ &= \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \sum_{j=0}^{\lfloor \frac{k-1}{s+1} \rfloor} (-1)^j \binom{n-pk}{j} \binom{n-pk+k-(s+1)j-2}{n-pk-1}. \end{aligned} \tag{2}$$

We now prove Theorem 1, which states that for  $1 \leq n \leq sp + 1$ ,

$$|\mathcal{A}_{p,n}^{(s)}| = \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \binom{n-pk+k-2}{k-1},$$

and for  $n \geq sp + 2$ , we have

$$|\mathcal{A}_{p,n}^{(s)}| = \sum_{i=0}^s |\mathcal{A}_{p,n-1-ip}^{(s)}|. \tag{3}$$

*Proof of Theorem 1.* We note that

$$\frac{k-1}{s+1} \leq \frac{\frac{sn+1}{sp+1} - 1}{s+1} \leq \frac{\frac{s(sp+1)+1}{sp+1} - 1}{s+1} = \frac{s + \frac{1}{sp+1} - 1}{s+1} < 1.$$

Hence, it follows from (2) that

$$\begin{aligned} |\mathcal{A}_{p,n}^{(s)}| &= \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \sum_{j=0}^{\lfloor \frac{k-1}{s+1} \rfloor} (-1)^j \binom{n-pk}{j} \binom{n-pk+k-(s+1)j-2}{n-pk-1} \\ &= \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \binom{n-pk+k-2}{n-pk-1} \\ &= \sum_{k=1}^{\lfloor \frac{sn+1}{sp+1} \rfloor} \binom{n-pk+k-2}{k-1}. \end{aligned}$$

To prove (3), we partition  $\mathcal{A}_{p,n}^{(s)}$  into  $(s+1)$  sets  $(\mathcal{A}_{p,n,i}^{(s)})_{i=0}^s$ , where

$$\mathcal{A}_{p,n,i}^{(s)} := \{F \in \mathcal{A}_{p,n}^{(s)} : F \text{ contains exactly } i \text{ copies of } (n-1)\}.$$

For each integer  $i \in [0, s]$ , define a map  $\psi_i : \mathcal{A}_{p,n-1-ip}^{(s)} \rightarrow \mathcal{A}_{p,n,i}^{(s)}$  as

$$F \longrightarrow \begin{cases} (F \setminus \{n-1\}) \cup \{n\}, & \text{if } i = 0; \\ \left( (F + ip) \cup \underbrace{\{n-1, \dots, n-1\}}_{i-1} \right) \cup \{n\}, & \text{if } 1 \leq i \leq s. \end{cases}$$

First, we show that each  $\psi_i$  is well-defined.

When  $i = 0$ , the map  $\psi_0$  increases the maximum element of the input set by 1, so if  $F \in \mathcal{A}_{p,n-1}^{(s)}$ , then  $\psi_0(F)$  does not contain  $(n-1)$ , contains exactly one copy of  $n$  as the maximum, and

$$\min \psi_0(F) \geq \min F \geq p|F| = p|\psi_0(F)|.$$

Hence,  $\psi_0(F) \in \mathcal{A}_{p,n,0}^{(s)}$ .

When  $1 \leq i \leq s$ , the map  $\psi_i$  increases the size of the input set by  $i$  and increases the minimum by  $ip$ , so

$$\min \psi_i(F) = \min F + ip \geq p|F| + ip = p|\psi_i(F)|.$$

Furthermore,  $\psi_i(F)$  contains exactly  $i$  copies of  $n-1$  and contains exactly one copy of  $n$  as the maximum. Hence,  $\psi_i(F) \in \mathcal{A}_{p,n,i}^{(s)}$ .

Next, we prove that each  $\psi_i$  is a bijection. Injectivity follows immediately from the definition of  $\psi_i$ . We show that each  $\psi_i$  is onto.

**Case 1:**  $G \in \mathcal{A}_{p,n,0}^{(s)}$ . Then  $n \in G$  but  $(n - 1) \notin G$ . Let  $F := (G \cup \{n - 1\}) \setminus \{n\}$ . We know that  $F$  contains exactly one copy of  $(n - 1)$  as the maximum, and

$$\min F = \begin{cases} \min G \geq p|G| = p|F|, & \text{if } |G| > 1; \\ n - 1 \geq p = p|F|, & \text{if } G = \{n\} \text{ (because } n \geq sp + 2). \end{cases}$$

Thus,  $F \in \mathcal{A}_{p,n-1}^{(s)}$  and  $\psi_0(F) = G$ .

**Case 2:**  $G \in \mathcal{A}_{p,n,i}^{(s)}$  with  $1 \leq i \leq s$ . Let

$$F := (G \setminus \underbrace{\{n, n - 1, \dots, n - 1\}}_{i-1}) - ip.$$

Since  $G$  has exactly  $i$  copies of  $(n - 1)$ , the set  $F$  contains exactly one copy of  $(n - 1 - ip)$  as the maximum. Furthermore,

$$\min F = \min G - ip \geq p|G| - ip = p(|G| - i) = p|F|.$$

Thus,  $F \in \mathcal{A}_{p,n-1-ip}^{(s)}$  and  $\psi_i(F) = G$ .

We have shown that  $|\mathcal{A}_{p,n-1-ip}^{(s)}| = |\mathcal{A}_{p,n,i}^{(s)}|$ . Therefore,

$$|\mathcal{A}_{p,n}^{(s)}| = \sum_{i=0}^s |\mathcal{A}_{p,n,i}^{(s)}| = \sum_{i=0}^s |\mathcal{A}_{p,n-1-ip}^{(s)}|.$$

□

### 3. Colored Generalized Schreier Multisets

We find a formula for  $|\mathcal{B}_{p,n}^{(s)}|$ . To form a  $k$ -element set, we choose  $(k - 1)$  elements in

$$\{(kp)_1, \dots, (kp)_s, \dots, (n - 1)_1, \dots, (n - 1)_s\}.$$

Hence, the number of  $k$ -element sets in  $\mathcal{B}_{p,n}^{(s)}$  is  $\binom{(n-kp)s}{k-1}$  with  $k - 1 \leq (n - kp)s$ , i.e.,  $k \leq (ns + 1)/(ps + 1)$ . Therefore,

$$|\mathcal{B}_{p,n}^{(s)}| = \sum_{k=1}^{\lfloor \frac{ns+1}{ps+1} \rfloor} \binom{(n-kp)s}{k-1}. \tag{4}$$

This proves the first statement of Theorem 2. We use characteristic polynomials to prove the second statement. The first step is to find a parent sequence, denoted

by  $(b_{p,n}^{(s)})_{n=1}^\infty$ , that has  $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^\infty$  as a periodic subsequence. The parent sequence  $(b_{p,n}^{(s)})_{n=1}^\infty$  should satisfy a relatively simple recurrence that we take advantage of later.

Given positive integers  $s$  and  $p$ , we define the sequence  $(b_{p,n}^{(s)})_{n=1}^\infty$  recursively as follows:

$$b_{p,1}^{(s)} = \dots = b_{p,(p-1)s}^{(s)} = 0, \quad b_{p,(p-1)s+1}^{(s)} = \dots = b_{p,ps+1}^{(s)} = 1, \quad \text{and}$$

$$b_{p,n}^{(s)} = b_{p,n-1}^{(s)} + b_{p,n-1-ps}^{(s)}, \quad \text{for } n \geq ps + 2.$$

For example,

$$(b_{1,n}^{(2)})_{n=1}^\infty \text{ (A000930)} : 1, 1, 1, 2, 3, 4, 6, 9, 13, 19, 28, 41, 60, 88, 129, 189, 277, 406, \dots;$$

$$(b_{2,n}^{(2)})_{n=1}^\infty \text{ (A003520)} : 0, 0, 1, 1, 1, 1, 1, 2, 3, 4, 5, 6, 8, 11, 15, 20, 26, 34, 45, 60, \dots;$$

$$(b_{2,n}^{(3)})_{n=1}^\infty \text{ (A005709)} : 0, 0, 0, 1, 1, 1, 1, 1, 1, 1, 2, 3, 4, 5, 6, 7, 8, 10, 13, 17, 22, 28, 35, \dots$$

**Proposition 1.** For  $(s, p, n) \in \mathbb{N}^3$ , it holds that

$$b_{p,n}^{(s)} = \sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i}.$$

*Proof.* For  $n \leq (p-1)s$ , we have

$$\frac{n-s(p-1)-1}{sp+1} \leq \frac{-1}{sp+1}, \text{ so } \left\lfloor \frac{n-s(p-1)-1}{sp+1} \right\rfloor \leq -1.$$

Hence,

$$\sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} = 0,$$

as desired.

For  $(p-1)s + 1 \leq n \leq ps + 1$ , we have

$$\frac{n-s(p-1)-1}{sp+1} \geq \frac{(p-1)s+1-s(p-1)-1}{sp+1} = 0$$

and

$$\frac{n-s(p-1)-1}{sp+1} \leq \frac{ps+1-s(p-1)-1}{sp+1} = \frac{s}{sp+1} < 1.$$

Thus,

$$\left\lfloor \frac{n-s(p-1)-1}{sp+1} \right\rfloor = 0,$$

which gives

$$\sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} = \binom{n-s(p-1)-1}{0} = 1.$$

For  $n \geq ps + 2$ , we show that

$$\begin{aligned} & \sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} \\ &= \sum_{i=0}^{\lfloor \frac{n-2-s(p-1)}{sp+1} \rfloor} \binom{n-s(p-1)-2-spi}{i} \\ &+ \sum_{i=0}^{\lfloor \frac{n-2-sp-s(p-1)}{sp+1} \rfloor} \binom{n-2-sp-s(p-1)-spi}{i}. \end{aligned}$$

We proceed by case analysis.

**Case 1:**  $n = m(sp + 1) + r$  with  $m \geq 1$  and  $1 \leq r \leq s(p - 1)$ . Then

$$\begin{aligned} & \sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} - \sum_{i=0}^{\lfloor \frac{n-2-s(p-1)}{sp+1} \rfloor} \binom{n-s(p-1)-2-spi}{i} \\ &= \sum_{i=0}^{m-1} \left( \binom{n-s(p-1)-1-spi}{i} - \binom{n-s(p-1)-2-spi}{i} \right) \\ &= \sum_{i=1}^{m-1} \left( \binom{n-s(p-1)-1-spi}{i} - \binom{n-s(p-1)-2-spi}{i} \right) \\ &= \sum_{i=1}^{m-1} \binom{n-s(p-1)-2-spi}{i-1} \\ &= \sum_{i=0}^{m-2} \binom{n-s(p-1)-2-sp(i+1)}{i} \\ &= \sum_{i=0}^{\lfloor \frac{n-2-sp-s(p-1)}{sp+1} \rfloor} \binom{n-2-sp-s(p-1)-spi}{i}, \end{aligned}$$

because

$$\left\lfloor \frac{n-2-sp-s(p-1)}{sp+1} \right\rfloor = \left\lfloor \frac{(m-2)(sp+1) + r + s}{sp+1} \right\rfloor = m-2.$$

**Case 2:**  $n = m(sp + 1) + s(p - 1) + 1$  with  $m \geq 1$ . We have

$$\begin{aligned}
 & \sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} - \sum_{i=0}^{\lfloor \frac{n-2-s(p-1)}{sp+1} \rfloor} \binom{n-s(p-1)-2-spi}{i} \\
 &= \sum_{i=0}^m \binom{n-s(p-1)-1-spi}{i} - \sum_{i=0}^{m-1} \binom{n-s(p-1)-2-spi}{i} \\
 &= 1 + \sum_{i=1}^{m-1} \left( \binom{n-s(p-1)-1-spi}{i} - \binom{n-s(p-1)-2-spi}{i} \right) \\
 &= 1 + \sum_{i=1}^{m-1} \binom{n-s(p-1)-2-spi}{i-1} \\
 &= 1 + \sum_{i=0}^{m-2} \binom{n-s(p-1)-2-sp(i+1)}{i} \\
 &= \sum_{i=0}^{m-1} \binom{n-s(p-1)-2-sp(i+1)}{i} \\
 &= \sum_{i=0}^{\lfloor \frac{n-2-sp-s(p-1)}{sp+1} \rfloor} \binom{n-2-sp-s(p-1)-spi}{i}.
 \end{aligned}$$

**Case 3:**  $n = m(sp + 1) + r$  with  $m \geq 1$  and  $s(p - 1) + 2 \leq r \leq sp$ . Then

$$\begin{aligned}
 & \sum_{i=0}^{\lfloor \frac{n-s(p-1)-1}{sp+1} \rfloor} \binom{n-s(p-1)-1-spi}{i} - \sum_{i=0}^{\lfloor \frac{n-2-s(p-1)}{sp+1} \rfloor} \binom{n-s(p-1)-2-spi}{i} \\
 &= \sum_{i=0}^m \left( \binom{n-s(p-1)-1-spi}{i} - \binom{n-s(p-1)-2-spi}{i} \right) \\
 &= \sum_{i=1}^m \left( \binom{n-s(p-1)-1-spi}{i} - \binom{n-s(p-1)-2-spi}{i} \right) \\
 &= \sum_{i=1}^m \binom{n-s(p-1)-2-spi}{i-1} \\
 &= \sum_{i=0}^{m-1} \binom{n-s(p-1)-2-sp(i+1)}{i} \\
 &= \sum_{i=0}^{\lfloor \frac{n-2-sp-s(p-1)}{sp+1} \rfloor} \binom{n-2-sp-s(p-1)-spi}{i}.
 \end{aligned}$$

□

**Lemma 1.** *The sequence  $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^\infty$  is a periodic subsequence of  $(b_{p,n}^{(s)})_{n=1}^\infty$ . In particular,*

$$|\mathcal{B}_{p,n}^{(s)}| = b_{p,ns-s+1}^{(s)}.$$

*Proof.* Thanks to (4) and Proposition 1, it suffices to verify that

$$\sum_{k=1}^{\lfloor \frac{ns+1}{ps+1} \rfloor} \binom{ns-sp k}{k-1} = \sum_{i=0}^{\lfloor \frac{(n-p)s}{ps+1} \rfloor} \binom{ns-sp(i+1)}{i}.$$

Indeed,

$$\sum_{k=1}^{\lfloor \frac{ns+1}{ps+1} \rfloor} \binom{ns-sp k}{k-1} = \sum_{i=0}^{\lfloor \frac{ns+1}{ps+1} \rfloor - 1} \binom{ns-sp(i+1)}{i} = \sum_{i=0}^{\lfloor \frac{(n-p)s}{ps+1} \rfloor} \binom{ns-sp(i+1)}{i}.$$

□

We now introduce characteristic polynomials that enable us to deduce a linear recurrence of a periodic subsequence from its parent sequence.

**Definition 1.** Let  $p(x) = c_0 + c_1x + \dots + c_r x^r$  be a polynomial with real coefficients  $(c_i)_{i=0}^r$ . A sequence  $(a_n)_{n=1}^\infty$  is said to satisfy  $p(x)$  if

$$c_0 a_n + c_1 a_{n-1} + \dots + c_r a_{n-r} = 0, \text{ for all } n \geq r + 1.$$

It is readily verified that given two polynomials  $p(x)$  and  $q(x)$  such that  $p(x)$  divides  $q(x)$ , if a sequence  $(a_n)_{n=1}^\infty$  satisfies  $p(x)$ , then  $(a_n)_{n=1}^\infty$  satisfies  $q(x)$ .

By definition, the sequence  $(b_{p,n}^{(s)})_{n=1}^\infty$  satisfies the polynomial

$$u_{s,p}(x) := 1 - x - x^{ps+1}.$$

The second statement of Theorem 2 says that  $(|\mathcal{B}_{p,n}^{(s)}|)_{n=1}^\infty$  satisfies

$$v_{s,p}(x) := 1 - \binom{s}{0}x - \binom{s}{1}x^{1+p} - \dots - \binom{s}{s}x^{1+sp} = 1 - \sum_{i=0}^s \binom{s}{i}x^{1+ip}.$$

*Proof of the second statement of Theorem 2.* We have

$$\begin{aligned} u_{s,p}(x) \cdot \sum_{i=0}^{s-1} x^i (x^{sp} + 1)^i &= (1 - x - x^{ps+1}) \frac{1 - x^s (x^{sp} + 1)^s}{1 - x (x^{sp} + 1)} \\ &= 1 - x^s (x^{sp} + 1)^s \\ &= 1 - x^s \sum_{i=0}^s \binom{s}{i} x^{sp i} \\ &= 1 - \sum_{i=0}^s \binom{s}{i} (x^s)^{1+ip} = v_{s,p}(x^s). \end{aligned}$$

Hence,  $u_{s,p}(x)$  divides  $v_{s,p}(x^s)$ , and so,  $(b_{p,n}^{(s)})_{n=1}^\infty$  satisfies  $v_{s,p}(x^s)$ . Due to the power  $s$  in  $v_{s,p}(x^s)$ , the polynomial  $v_{s,p}(x^s)$  gives a linear recurrence for terms in  $(b_{p,n}^{(s)})_{n=1}^\infty$  whose indices are  $s$  apart. It follows from Lemma 1 that the sequence  $(|\mathcal{B}_{p,n}|)_{n=1}^\infty$  satisfies  $v_{s,p}(x)$ .  $\square$

#### 4. Schreier Sets That Do Not Admit Multiples of a Given Number

In this section, given  $u \geq 2$ , we count Schreier sets that do not contain any multiple of  $u$ . This is the opposite of what was studied in [9]. First, we need a formula that gives all positive integers that are not divisible by  $u$ .

**Lemma 2.** *For  $u \geq 2$ , we have*

$$\left\{ \left\lfloor \frac{un - 1}{u - 1} \right\rfloor : n \in \mathbb{N} \right\} = \{n \in \mathbb{N} : u \nmid n\}.$$

*Proof.* Write  $n = (u - 1)j + m$  with  $j \geq 0$  and  $0 \leq m \leq u - 2$ . Then

$$\begin{aligned} \left\lfloor \frac{un - 1}{u - 1} \right\rfloor &= \left\lfloor \frac{u((u - 1)j + m) - 1}{u - 1} \right\rfloor = uj + m + \left\lfloor \frac{m - 1}{u - 1} \right\rfloor \\ &= \begin{cases} uj - 1, & \text{if } m = 0; \\ uj + m, & \text{if } 1 \leq m \leq u - 2. \end{cases} \end{aligned}$$

This shows that the formula  $\lfloor (un - 1)/(u - 1) \rfloor$  gives all positive integers that are not divisible by  $u$ .  $\square$

We prove Theorem 3 using the same method of characteristic polynomials as in Section 3. First, we find a formula for  $|\mathcal{D}_{u,n}|$ . A  $k$ -element set in  $\mathcal{D}_{u,n}$  is uniquely determined by  $(k - 1)$  elements in

$$\left\{ \left\lfloor \frac{u(j_k - 1) - 1}{u - 1} \right\rfloor, \dots, \left\lfloor \frac{u(n - 1) - 1}{u - 1} \right\rfloor \right\},$$

where  $j_k$  is the smallest positive integer with

$$\frac{u(j_k - 1) - 1}{u - 1} \geq k,$$

which is equivalent to

$$j_k = k + 1 - \left\lfloor \frac{k - 1}{u} \right\rfloor.$$

Hence, the number of  $k$ -element sets in  $\mathcal{D}_{u,n}$  is

$$\binom{n - j_k + 1}{k - 1} = \binom{n - k + \lfloor \frac{k-1}{u} \rfloor}{k - 1}$$

under the condition

$$n - k + \left\lfloor \frac{k-1}{u} \right\rfloor \geq k - 1; \text{ that is, } k \leq \frac{(n+1)u - 1}{2u - 1}.$$

Therefore,

$$\begin{aligned} |\mathcal{D}_{u,n}| &= \sum_{k=1}^{\lfloor \frac{(n+1)u-1}{2u-1} \rfloor} \binom{n-k + \lfloor \frac{k-1}{u} \rfloor}{k-1} = \sum_{k=0}^{\lfloor \frac{nu-u}{2u-1} \rfloor} \binom{n-k-1 + \lfloor \frac{k}{u} \rfloor}{k} \\ &= \sum_{k=0}^{\lfloor \frac{nu-u}{2u-1} \rfloor} \binom{\lfloor \frac{un-(u-1)k}{u} \rfloor - 1}{k}. \end{aligned} \tag{5}$$

We now define parent sequences  $(d_{u,n})_{n=1}^\infty$  so that for each fixed  $u$ , the sequence  $(|\mathcal{D}_{u,n}|)_{n=1}^\infty$  is a periodic subsequence of  $(d_{u,n})_{n=1}^\infty$ . Let

$$d_{u,1} = d_{u,2} = \dots = d_{u,2u-1} = 1$$

and

$$d_{u,n} = d_{u,n-u} + d_{u,n-2u+1} \text{ for } n \geq 2u.$$

For example,

- $(d_{2,n})_{n=1}^\infty : 1, 1, 1, 2, 2, 3, 4, 5, 7, 9, 12, 16, 21, 28, 37, 49, 65, 86, 114, 151, 200, 265, 351, \dots;$
- $(d_{3,n})_{n=1}^\infty : 1, 1, 1, 1, 1, 2, 2, 2, 3, 3, 4, 5, 5, 7, 8, 9, 12, 13, 16, 20, 22, 28, 33, 38, 48, 55, 66, 81, 93, 114, 136, 159, 195, 229, \dots;$
- $(d_{4,n})_{n=1}^\infty : 1, 1, 1, 1, 1, 1, 1, 2, 2, 2, 2, 3, 3, 3, 4, 5, 5, 5, 7, 8, 8, 9, 12, 13, 13, 16, 20, 21, 22, 28, 33, 34, 38, 48, 54, 56, 66, 81, 88, 94, 114, 135, 144, 160, 195, \dots$

**Proposition 2.** For  $u \geq 2$  and  $n \in \mathbb{N}$ , we have

$$d_{u,n} = \sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i}.$$

Note that when  $n < u$ , the sum consists of only  $\binom{-1}{0}$ , which is 1.

*Proof.* Let  $u \geq 2$ . If  $1 \leq n \leq u$ , then  $\lfloor \frac{n-1}{u} \rfloor = 0$ , and

$$\sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} = \binom{\lfloor \frac{n-u}{2u-1} \rfloor}{0} = 1.$$

If  $u + 1 \leq n \leq 2u - 1$ , then

$$1 \leq \frac{n-1}{u} \leq \frac{2u-2}{u} = 2 - \frac{2}{u} < 2 \implies \left\lfloor \frac{n-1}{u} \right\rfloor = 1,$$

and

$$\begin{aligned} \sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} &= \sum_{i=0}^1 \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} \\ &= \binom{\lfloor \frac{n-u}{2u-1} \rfloor}{0} + \binom{\lfloor \frac{n-1}{2u-1} \rfloor}{1} \\ &= \binom{0}{0} + \binom{0}{1} = 1. \end{aligned}$$

It remains to verify that  $d_{u,n} = d_{u,n-u} + d_{u,n-2u+1}$  for all  $n \geq 2u$ , i.e.,

$$\sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} = \sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor - 1} \binom{\lfloor \frac{n+(u-1)i-2u}{2u-1} \rfloor}{i} + \sum_{i=0}^{\lfloor \frac{n}{u} \rfloor - 2} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor - 1}{i}.$$

By Pascal's rule, we have

$$\begin{aligned} &\sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} \\ &= 1 + \sum_{i=1}^{\lfloor \frac{n-1}{u} \rfloor} \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} \\ &= 1 + \underbrace{\sum_{i=1}^{\lfloor \frac{n-1}{u} \rfloor} \left( \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} - 1 \right)}_{=: I(u,n)} + \underbrace{\sum_{i=1}^{\lfloor \frac{n-1}{u} \rfloor} \left( \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i-1} - 1 \right)}_{=: II(u,n)}. \end{aligned}$$

Reindexing  $II(u, n)$  gives

$$II(u, n) = \sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor - 1} \binom{\lfloor \frac{n+(u-1)(i+1)-u-2u+1}{2u-1} \rfloor}{i} = \sum_{i=0}^{\lfloor \frac{n-1}{u} \rfloor - 1} \binom{\lfloor \frac{n+(u-1)i-2u}{2u-1} \rfloor}{i}.$$

Hence, it suffices to show that

$$I(u, n) = \sum_{i=1}^{\lfloor \frac{n}{u} \rfloor - 2} \left( \binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor}{i} - 1 \right),$$

which means that

$$\binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor - 1}{i} = 0, \text{ for } \lfloor \frac{n}{u} \rfloor - 1 \leq i \leq \lfloor \frac{n-1}{u} \rfloor.$$

Write  $n = qu + r$ , where  $q \geq 2$  and  $0 \leq r \leq u - 1$ . Then

$$\lfloor \frac{n}{u} \rfloor - 1 = q - 1 \quad \text{and} \quad \lfloor \frac{n-1}{u} \rfloor = \begin{cases} q, & \text{if } r \geq 1; \\ q - 1, & \text{if } r = 0. \end{cases}$$

**Case 1:** if  $i = q - 1$ , then

$$\lfloor \frac{n + (u-1)(q-1) - u}{2u-1} \rfloor = \lfloor \frac{(2u-1)q + r - (2u-1)}{2u-1} \rfloor = q - 1.$$

Hence,

$$\binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor - 1}{q-1} = 0.$$

**Case 2:** if  $i = q$ , then

$$\lfloor \frac{n + (u-1)q - u}{2u-1} \rfloor = \lfloor \frac{(2u-1)q + r - u}{2u-1} \rfloor = q - 1.$$

Thus,

$$\binom{\lfloor \frac{n+(u-1)i-u}{2u-1} \rfloor - 1}{q} = 0.$$

This completes our proof. □

**Proposition 3.** For each  $u \geq 2$ , the sequence  $(|\mathcal{D}_{u,n}|)_{n=1}^\infty$  is a periodic subsequence of  $(d_{u,n})_{n=1}^\infty$ . In particular,

$$|\mathcal{D}_{u,n}| = d_{u,un-(u-1)}, \text{ for all } n \in \mathbb{N}.$$

*Proof.* Fix  $u \geq 2$  and  $n \in \mathbb{N}$ . By Proposition 2,

$$d_{u,un-(u-1)} = \sum_{i=0}^{\lfloor \frac{un-u+1-1}{u} \rfloor} \binom{\lfloor \frac{un-u+1+(u-1)i-u}{2u-1} \rfloor}{i} = \sum_{j=0}^{n-1} \binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j}.$$

Thanks to (5), it suffices to prove that

$$\sum_{k=0}^{\lfloor \frac{nu-u}{2u-1} \rfloor} \underbrace{\binom{\lfloor \frac{un-(u-1)k}{u} \rfloor - 1}{k}}_{=:f(k)} = \sum_{j=0}^{n-1} \underbrace{\binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j}}_{=:g(j)}. \tag{6}$$

If  $n = 1$ , both sides of (6) equal 1. Suppose that  $n \geq 2$ . For  $0 \leq j \leq n - 1$ , we consider  $j$  of the form

$$j = n - (2u - 1)(\ell + 1) =: \xi(\ell).$$

Then

$$0 \leq \ell \leq \underbrace{\left\lfloor \frac{n - 2u}{2u - 1} + \frac{1}{2u - 1} \right\rfloor}_{=: \ell_0} < \frac{n - 2u}{2u - 1} + \frac{1}{u}.$$

For  $\ell$  in the above range, if  $k = \chi(\ell) := u(\ell + 1) - 1$ , then  $0 \leq k \leq \left\lfloor \frac{nu - u}{2u - 1} \right\rfloor$ . We have

$$\begin{aligned} g(\xi(\ell)) &= g(n - (2u - 1)(\ell + 1)) \\ &= \binom{\left\lfloor \frac{un + (u-1)(n - (2u-1)(\ell+1))}{2u-1} \right\rfloor - 1}{n - (2u - 1)(\ell + 1)} \\ &= \binom{\left\lfloor \frac{(2u-1)n - (u-1)(2u-1)(\ell+1)}{2u-1} \right\rfloor - 1}{n - (2u - 1)(\ell + 1)} \\ &= \binom{n + \ell - (\ell + 1)u}{n - (2u - 1)(\ell + 1)} \\ &= \binom{n + \ell - (\ell + 1)u}{n + \ell - (\ell + 1)u - (n - (2u - 1)(\ell + 1))} \\ &= \binom{n + \ell - (\ell + 1)u}{(\ell + 1)u - 1} \\ &= \binom{\left\lfloor n - (u - 1)(\ell + 1) - \frac{1}{u} \right\rfloor}{u\ell + u - 1} \\ &= \binom{\left\lfloor \frac{un - (u-1)(u\ell + u - 1)}{u} \right\rfloor - 1}{u\ell + u - 1} \\ &= f(u\ell + (u - 1)) = f(\chi(\ell)). \end{aligned} \tag{7}$$

Write

$$\begin{aligned} \sum_{k=0}^{\left\lfloor \frac{nu - u}{2u - 1} \right\rfloor} f(k) &= \sum_{k=0}^{\chi(0)-1} f(k) + f(\chi(0)) + \sum_{k=\chi(0)+1}^{\chi(1)-1} f(k) + f(\chi(1)) + \dots \\ &\quad + \sum_{k=\chi(\ell_0-1)+1}^{\chi(\ell_0)-1} f(k) + f(\chi(\ell_0)) + \sum_{k=\chi(\ell_0)+1}^{\left\lfloor \frac{nu - u}{2u - 1} \right\rfloor} f(k) \end{aligned}$$

and

$$\begin{aligned} \sum_{j=0}^{n-1} g(j) &= \sum_{j=0}^{\xi(\ell_0)-1} g(j) + g(\xi(\ell_0)) + \sum_{j=\xi(\ell_0)+1}^{\xi(\ell_0-1)-1} g(j) + g(\xi(\ell_0 - 1)) + \dots \\ &\quad + \sum_{j=\xi(1)+1}^{\xi(0)-1} g(j) + g(\xi(0)) + \sum_{j=\xi(0)+1}^{n-1} g(j). \end{aligned}$$

Note that (7) gives  $f(\chi(\ell)) = g(\xi(\ell))$  for each  $\ell \in [0, \ell_0]$ .

Next, we prove that for  $\ell \in [0, \ell_0 - 1]$ ,

$$\sum_{k=\chi(\ell)+1}^{\chi(\ell+1)-1} f(k) = \sum_{j=\xi(\ell+1)+1}^{\xi(\ell)-1} g(j). \tag{8}$$

To do so, note that the right sum of (8) has  $2u - 2$  terms, while the left sum has  $u - 1$  terms. To prove (8), we use Pascal’s rule to combine every two consecutive terms in the right sum. On the one hand, we have

$$\begin{aligned} &\sum_{j=\xi(\ell+1)+1}^{\xi(\ell)-1} g(j) \\ &= \sum_{i=1}^{u-1} (g(\xi(\ell + 1) + 2i - 1) + g(\xi(\ell + 1) + 2i)) \\ &= \sum_{i=1}^{u-1} \left( \binom{n - (u - 1)(\ell + 2) + i - 2}{\xi(\ell + 1) + 2i - 1} + \binom{n - (u - 1)(\ell + 2) + i - 2}{\xi(\ell + 1) + 2i} \right) \\ &= \sum_{i=1}^{u-1} \binom{n - (u - 1)(\ell + 2) + i - 1}{\xi(\ell + 1) + 2i} = \sum_{i=1}^{u-1} \binom{n - (u - 1)(\ell + 2) + i - 1}{u(\ell + 2) - i - 1}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \sum_{k=\chi(\ell)+1}^{\chi(\ell+1)-1} f(k) &= \sum_{i=1}^{u-1} f(\chi(\ell + 1) - i) \\ &= \sum_{i=1}^{u-1} \binom{\lfloor \frac{un - (u-1)(\chi(\ell+1) - i)}{u} \rfloor - 1}{\chi(\ell + 1) - i} \\ &= \sum_{i=1}^{u-1} \binom{n - (u - 1)(\ell + 2) + i - 1}{u(\ell + 2) - i - 1}. \end{aligned}$$

Therefore, (8) holds.

From (7) and (8), we have shown that

$$\sum_{k=\chi(0)}^{\chi(\ell_0)} f(k) = \sum_{j=\xi(\ell_0)}^{\xi(0)} g(j).$$

It remains to verify that

$$\sum_{k=0}^{\chi(0)-1} f(k) = \sum_{j=\xi(0)+1}^{n-1} g(j) \tag{9}$$

and

$$\sum_{k=\chi(\ell_0)+1}^{\lfloor \frac{n\chi(\ell_0)-1}{2\chi(\ell_0)-1} \rfloor} f(k) = \sum_{j=0}^{\xi(\ell_0)-1} g(j). \tag{10}$$

The proofs of (9) and (10) use a similar idea to the proof of (8), so we move them to the Appendix.  $\square$

Theorem 3 shows how to obtain the sequence  $(|\mathcal{D}_{u,n}|)_{n=1}^\infty$  recursively: for  $1 \leq n \leq 2u - 1$ , we have  $|\mathcal{D}_{u,n}| = F_n$ , and for  $n \geq 2u$ ,

$$|\mathcal{D}_{u,n}| = \sum_{i=1}^u (-1)^{i-1} \binom{u}{i} |\mathcal{D}_{u,n-i}| + |\mathcal{D}_{u,n-2u+1}|. \tag{11}$$

*Proof of Theorem 3.* The first statement follows from the well-known formula for  $F_n$ :

$$F_n = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-i-1}{i}. \tag{12}$$

For a proof of (12) using tilings, see [4, Identity 4], or [10, (2.5)] gives another proof via Zeckendorf’s Theorem [12].

Combining (5) with (12), we need to show the equality

$$\sum_{k=1}^{\lfloor \frac{(n+1)u-1}{2u-1} \rfloor} \binom{n-k + \lfloor \frac{k-1}{u} \rfloor}{k-1} = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-i-1}{i} \text{ for } n \leq 2u - 1, \tag{13}$$

whose proof is in the Appendix.

By definition, the sequence  $(d_{u,n})_{n=1}^\infty$  satisfies the polynomial

$$p_u(x) := 1 - x^u - x^{2u-1}.$$

Recurrence (11) says that  $(|\mathcal{D}_{k,n}|)_{n=1}^\infty$  satisfies

$$q_u(x) := 1 + \sum_{i=1}^u \binom{u}{i} (-x)^i - x^{2u-1} = (1-x)^u - x^{2u-1}.$$

By Lemma 3.4 and Proposition 3, it suffices to verify that  $p_u(x)$  divides  $q_u(x^u)$ . We have

$$\begin{aligned} & (1 - x^u - x^{2u-1}) \left( \sum_{j=0}^{u-1} (1 - x^u)^{u-1-j} x^{j(2u-1)} \right) \\ &= (1 - x^u - x^{2u-1}) \frac{(1 - x^u)^u - x^{u(2u-1)}}{1 - x^u - x^{2u-1}} \\ &= (1 - x^u)^u - x^{u(2u-1)} = q_u(x^u). \end{aligned}$$

□

### 5. Further Investigations

One may consider other setups for  $p, q$ , and  $(s_{n,i})_{i=1}^n$  in Problem 1 and determine what recurrences they give. Interested readers may also study the more general condition  $q \min F \geq p|F|$  as in [2]. Define

$$\mathcal{A}_{p,q,n}^{(s)} := \{F \subset \underbrace{\{1, \dots, 1\}}_s, \dots, \underbrace{\{n-1, \dots, n-1\}}_s, n\} : n \in F \text{ and } q \min F \geq p|F|\}.$$

Similar to (2), we have a formula for  $|\mathcal{A}_{n,p,q}^{(s)}|$ :

$$|\mathcal{A}_{p,q,n}^{(s)}| = \sum_{k=1}^{\lfloor \frac{nsq+q}{sp+q} \rfloor} \sum_{j=0}^{\lfloor \frac{k-1}{s+1} \rfloor} (-1)^j \binom{n - \lceil \frac{pk}{q} \rceil}{j} \binom{n - \lceil \frac{pk}{q} \rceil + k - 1 - (s+1)j - 1}{n - \lceil \frac{pk}{q} \rceil - 1}.$$

Below are the data and conjectured recurrences we gather:

1.  $(|\mathcal{A}_{1,2,n}^{(2)}|)_{n=1}^\infty$ : 1, 2, 4, 9, 19, 41, 88, 189, 406, 872, 1873, 4023, 8641, ... with

$$a_n = a_{n-1} + 2a_{n-2} + a_{n-3};$$

2.  $(|\mathcal{A}_{1,3,n}^{(2)}|)_{n=1}^\infty$ : 1, 3, 6, 13, 31, 73, 169, 392, 912, 2121, 4930, 11460, 26642, ... with

$$a_n = 3a_{n-1} - 3a_{n-2} + 4a_{n-3} - 2a_{n-4} + a_{n-5};$$

3.  $(|\mathcal{A}_{1,4,n}^{(2)}|)_{n=1}^\infty$ :

1, 3, 8, 18, 41, 100, 250, 617, 1501, 3643, 8877, 21689, 52984, 129303, ... with

$$a_n = 3a_{n-1} - 3a_{n-2} + 3a_{n-3} + 2a_{n-4} + a_{n-5};$$

4.  $(|\mathcal{A}_{1,5,n}^{(2)}|)_{n=1}^\infty$ :

1, 3, 9, 23, 54, 127, 314, 808, 2090, 5326, 13379, 33460, 83979, 211847, ... with

$$a_n = 5a_{n-1} - 10a_{n-2} + 10a_{n-3} - 4a_{n-5} + a_{n-6} + a_{n-7};$$

5.  $(|\mathcal{A}_{3,2,n}^{(2)}|)_{n=1}^\infty$ : 0, 1, 1, 2, 3, 5, 8, 14, 24, 40, 66, 110, 185, 311, 521, 871, ... with

$$a_n = a_{n-1} + 2a_{n-4} + a_{n-5} + a_{n-6} + a_{n-7}.$$

**Problem 2.** For  $(s, p, q) \in \mathbb{N}^3$ , find the recurrence for  $(|\mathcal{A}_{p,q,n}^{(s)}|)_{n=1}^\infty$ .

Observe that [2, Theorem 1.1] and Theorem 1 solve Problem 2 when  $s = 1$  and  $q = 1$ , respectively.

Lastly, we are interested in nonlinear Schreier-type conditions  $(\min F)^q \geq |F|^p$  as started in [7].

**Acknowledgements.** The authors thank the anonymous referee for the careful reading of the paper and for the valuable suggestions. This work was partially supported by the National Science Foundation DMS2341670. We thank the participants at Polymath Jr. 2025 for helpful discussions.

**References**

[1] K. Beanland and H. V. Chu, On Schreier-type sets, partitions, and compositions, *J. Integer Seq.* **27** (2024), 1–13.

[2] K. Beanland, H. V. Chu, and C. E. Finch-Smith, Generalized Schreier sets, linear recurrence relation, and Turán graphs, *Fibonacci Quart.* **60** (2022), 352–356.

[3] K. Beanland, D. Gorovoy, J. Hodor, and D. Homza, Counting unions of Schreier sets, *Bull. Aust. Math. Soc.* **110** (2024), 19–31.

[4] A. T. Benjamin and J. J. Quinn, *Proofs That Really Count: The Art of Combinatorial Proof*, Mathematical Association of America, Washington, D.C., 2003.

[5] A. Bird, Jozef Schreier, Schreier sets, and the Fibonacci sequence, blog post, <https://outofthenormmaths.wordpress.com/2012/05/13/jozef-schreier-schreier-sets-and-the-fibonacci-sequence/>.

[6] B. A. Bondarenko, Generalized Pascal triangles and generalized binomial coefficients, in *Generalized Pascal Triangles and Pyramids, Their Fractals, Graphs, and Applications*, The Fibonacci Association, 1993, 15–24.

[7] H. V. Chu, N. Irmak, S. J. Miller, L. Szalay, and S. X. Zhang, Schreier multisets and the  $s$ -step Fibonacci sequences, *Integers* **24A** (2024), 1–11.

[8] H. V. Chu, S. J. Miller, and Z. Xiang, Higher order Fibonacci sequences from generalized Schreier sets, *Fibonacci Quart.* **58** (2020), 249–253.

- [9] H. V. Chu and Z. L. Vasseur, Linear recurrences of generalized Schreier sets revisited, *J. Integer Seq.* **29** (2026), 1–18.
- [10] M. Koloğlu, G. S. Kopp, S. J. Miller, and Y. Wang, On the number of summands in Zeckendorf decompositions, *Fibonacci Quart.* **49** (2011), 116–130.
- [11] OEIS Foundation Inc., The On-Line Encyclopedia of Integer Sequences, <https://oeis.org>.
- [12] E. Zeckendorf, Representation des nombres naturels par une somme de nombres de Fibonacci ou denombres de Lucas, *Bull. Soc. Roy. Sci. Liege* **41** (1972), 179–182.

**Appendix: Technical Proofs**

*Proof of (9) that*  $\sum_{k=0}^{u-2} \binom{\lfloor \frac{un-(u-1)k}{u} \rfloor - 1}{k} = \sum_{j=n-2u+2}^{n-1} \binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j}$ . We have

$$\sum_{k=0}^{u-2} \binom{\lfloor \frac{un-(u-1)k}{u} \rfloor - 1}{k} = \sum_{k=0}^{u-2} \binom{n-k-1 + \lfloor \frac{k}{u} \rfloor}{k} = \sum_{k=0}^{u-2} \binom{n-k-1}{k},$$

while

$$\begin{aligned} & \sum_{j=n-2u+2}^{n-1} \binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j} \\ &= \sum_{j=n-2u+2}^{n-3} \binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j} + 1 \\ &= \sum_{i=1}^{u-2} \left( \binom{\lfloor \frac{un+(u-1)(n-2u+2i)}{2u-1} \rfloor - 1}{n-2u+2i} + \binom{\lfloor \frac{un+(u-1)(n-2u+2i+1)}{2u-1} \rfloor - 1}{n-2u+2i+1} \right) + 1 \\ &= \sum_{i=1}^{u-2} \left( \binom{n-u+i-1 + \lfloor \frac{u-i}{2u-1} \rfloor}{n-2u+2i} + \binom{n-u+i + \lfloor \frac{-i}{2u-1} \rfloor}{n-2u+2i+1} \right) + 1 \\ &= \sum_{i=1}^{u-2} \left( \binom{n-u+i-1}{n-2u+2i} + \binom{n-u+i-1}{n-2u+2i+1} \right) + 1 \\ &= \sum_{i=1}^{u-2} \binom{n-u+i}{n-2u+2i+1} + 1 \\ &= \sum_{i=1}^{u-2} \binom{n-u+i}{u-i-1} + 1 \\ &= \sum_{k=0}^{u-2} \binom{n-k-1}{k} \text{ by setting } u = k + i + 1. \end{aligned}$$

Hence, (9) holds. □

Regarding (10), we have

$$\xi(\ell_0) - 1 = n - 1 - (2u - 1) \left\lfloor \frac{n}{2u - 1} \right\rfloor \text{ and } \chi(\ell_0) + 1 = u \left\lfloor \frac{n}{2u - 1} \right\rfloor.$$

Hence, (10) is the same as

$$\sum_{k=u}^{\lfloor \frac{nu-u}{2u-1} \rfloor} \binom{\lfloor \frac{un-(u-1)k}{u} \rfloor - 1}{k} = \sum_{j=0}^{n-1-(2u-1)\lfloor \frac{n}{2u-1} \rfloor} \binom{\lfloor \frac{un+(u-1)j}{2u-1} \rfloor - 1}{j}. \tag{14}$$

*Proof of (10).* Write  $n = (2u - 1)s + t$  with  $s \geq 0$  and  $0 \leq t \leq 2u - 2$ . If  $t = 0$ , both sides of (14) equal 0. Assume that  $1 \leq t \leq 2u - 2$  and rewrite (14) as

$$\sum_{k=us}^{us+\lfloor \frac{(t-1)u}{2u-1} \rfloor} \binom{2us+t-k-1}{k} = \sum_{j=0}^{t-1} \binom{su+\lfloor \frac{tu+(u-1)j}{2u-1} \rfloor - 1}{j}.$$

**Case 1:**  $t = 2r + 1$  for some  $r \geq 0$ . Then  $r \leq u - 2$ . We have

$$\begin{aligned} & \sum_{j=0}^{t-1} \binom{su+\lfloor \frac{tu+(u-1)j}{2u-1} \rfloor - 1}{j} \\ &= \sum_{j=0}^{2r} \binom{su+\lfloor \frac{(2r+1)u+(u-1)j}{2u-1} \rfloor - 1}{j} \\ &= 1 + \sum_{i=1}^r \left( \binom{su+\lfloor \frac{(2r+1)u+(u-1)(2i-1)}{2u-1} \rfloor - 1}{2i-1} + \binom{su+\lfloor \frac{(2r+1)u+(u-1)2i}{2u-1} \rfloor - 1}{2i} \right) \\ &= 1 + \sum_{i=1}^r \left( \binom{su+r+i+\lfloor \frac{r-i+1}{2u-1} \rfloor - 1}{2i-1} + \binom{su+r+i+\lfloor \frac{r-i+u}{2u-1} \rfloor - 1}{2i} \right) \\ &= 1 + \sum_{i=1}^r \left( \binom{su+r+i-1}{2i-1} + \binom{su+r+i-1}{2i} \right) \\ &= 1 + \sum_{i=1}^r \binom{su+r+i}{2i} \end{aligned}$$

and

$$\begin{aligned} \sum_{k=us}^{us+\lfloor \frac{(t-1)u}{2u-1} \rfloor} \binom{2us+t-k-1}{k} &= \sum_{k=us}^{us+r} \binom{2us+2r-k}{k} \\ &= \sum_{k=us}^{us+r} \binom{2us+2r-k}{2us+2r-2k} \\ &= \sum_{j=0}^r \binom{us+r+j}{2j} \text{ by setting } j = r - k + us. \end{aligned}$$

Hence, (14) holds in this case.

Case 2:  $t = 2r$  for  $r \geq 1$ . Then  $r \leq u - 1$ . We have

$$\begin{aligned} & \sum_{j=0}^{t-1} \binom{su + \lfloor \frac{2ru+(u-1)j}{2u-1} \rfloor - 1}{j} \\ &= \sum_{j=0}^{2r-1} \binom{su + \lfloor \frac{2ru+(u-1)j}{2u-1} \rfloor - 1}{j} \\ &= \sum_{i=0}^{r-1} \left( \binom{su + \lfloor \frac{2ru+(u-1)2i}{2u-1} \rfloor - 1}{2i} + \binom{su + \lfloor \frac{2ru+(u-1)(2i+1)}{2u-1} \rfloor - 1}{2i+1} \right) \\ &= \sum_{i=0}^{r-1} \left( \binom{su+r+i + \lfloor \frac{r-i}{2u-1} \rfloor - 1}{2i} + \binom{su+r+i + \lfloor \frac{r+u-i-1}{2u-1} \rfloor - 1}{2i+1} \right) \\ &= \sum_{i=0}^{r-1} \left( \binom{su+r+i-1}{2i} + \binom{su+r+i-1}{2i+1} \right) \\ &= \sum_{i=0}^{r-1} \binom{su+r+i}{2i+1} \end{aligned}$$

and

$$\begin{aligned} \sum_{k=us}^{us+\lfloor \frac{(t-1)u}{2u-1} \rfloor} \binom{2us+t-k-1}{k} &= \sum_{k=us}^{us+r-1} \binom{2us+2r-k-1}{k} \\ &= \sum_{k=us}^{us+r-1} \binom{2us+2r-k-1}{2us+2r-2k-1} \\ &= \sum_{j=0}^{r-1} \binom{us+r+j}{2j+1} \text{ by setting } j = us+r-k-1. \end{aligned}$$

Thus, (14) holds in this case as well. □

We prove (13), which states that for  $n \leq 2u - 1$ ,

$$\sum_{k=1}^{\lfloor \frac{(n+1)u-1}{2u-1} \rfloor} \binom{n-k + \lfloor \frac{k-1}{u} \rfloor}{k-1} = \sum_{i=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n-i-1}{i}.$$

*Proof of (13).* Since  $n \leq 2u - 1$ ,

$$\frac{k-1}{u} \leq \frac{\frac{(n+1)u-1}{2u-1} - 1}{u} = \frac{(n-1)u}{u(2u-1)} = \frac{n-1}{2u-1} \leq \frac{2u-2}{2u-1},$$

so  $\lfloor \frac{k-1}{u} \rfloor = 0$ . Hence,

$$\sum_{k=1}^{\lfloor \frac{(n+1)u-1}{2u-1} \rfloor} \binom{n-k + \lfloor \frac{k-1}{u} \rfloor}{k-1} = \sum_{k=1}^{\lfloor \frac{(n+1)u-1}{2u-1} \rfloor} \binom{n-k}{k-1} = \sum_{j=0}^{\lfloor \frac{(n-1)u}{2u-1} \rfloor} \binom{n-j-1}{j}.$$

It remains to verify that

$$\left\lfloor \frac{(n-1)u}{2u-1} \right\rfloor = \left\lfloor \frac{n-1}{2} \right\rfloor, \text{ for } 1 \leq n \leq 2u-1. \tag{15}$$

If  $n = 2r - 1$  with  $1 \leq r \leq u$ , then

$$\frac{(n-1)u}{2u-1} = \frac{2u}{2u-1}(r-1) \in [r-1, r).$$

If  $n = 2r$  with  $1 \leq r \leq u-1$ , then

$$\frac{(n-1)u}{2u-1} = \frac{(2r-1)u}{2u-1} \in [r-1, r).$$

In both cases, we have (15). □