



NOTES ON SUM-FREE SETS IN ABELIAN GROUPS

Nathanaël Hassler

LIB, Université Bourgogne Europe, Dijon, France
 nathanael.hassler@ens-rennes.fr

Andrew Treglown

University of Birmingham, Birmingham, United Kingdom
 a.c.treglown@bham.ac.uk

Received: 6/20/25, Accepted: 7/13/26, Published: 7/29/26

Abstract

In this paper we highlight a few open problems concerning maximal sum-free sets in abelian groups. In addition, for most even order abelian groups G we asymptotically determine the number of maximal distinct sum-free subsets in G . Our proof makes use of the container method.

1. Introduction

Let $(G, +)$ be an abelian group. A triple $x, y, z \in G$ is a *Schur triple* if $x + y = z$; if additionally x, y, z are distinct, we call them a *distinct Schur triple*. A subset $S \subseteq G$ is a *sum-free set* if S does not contain any Schur triple. Similarly, we say S is a *distinct sum-free set* if S does not contain any distinct Schur triple. A sum-free set $S \subseteq G$ is *maximal* if S is not properly contained in another sum-free subset of G ; we define the notion of a *maximal distinct sum-free subset* analogously. We let $\mu(G)$ denote the size of the largest sum-free subset of G and $\mu^*(G)$ denote the size of the largest distinct sum-free subset of G .

The study of $\mu(G)$ dates back to the 1960s [10] and we now (through work of Green and Ruzsa [12]) have a complete understanding of the exact value of $\mu(G)$ for all abelian groups. To articulate this behavior we need the following definition.

Definition 1. Let G be an abelian group of order n .

- Let n be divisible by a prime $p \equiv 2 \pmod{3}$. Given the smallest such p , we say that G is *Type I(p)*.
- If n is not divisible by any prime $p \equiv 2 \pmod{3}$, but $3|n$, then we say that G is *Type II*.
- Otherwise, G is *Type III*.

DOI: 10.5281/zenodo.21678439

Theorem 1 ([10, 12]). *Given any finite abelian group G , if G is Type I(p) then $\mu(G) = |G| \left(\frac{1}{3} + \frac{1}{3p} \right)$. Otherwise, if G is Type II then $\mu(G) = \frac{|G|}{3}$. Finally, if G is Type III then $\mu(G) = |G| \left(\frac{1}{3} - \frac{1}{3m} \right)$ where m is the exponent (largest order of any element) of G .*

The case of Type I and II groups in Theorem 1 is due to Diananda and Yap [10]; the case of Type III groups is due to Green and Ruzsa [12]. Notice that Theorem 1 tells us that for every finite abelian group G of order n , we have that $2n/7 \leq \mu(G) \leq n/2$.

In recent years there has been significant interest in the study of transferring (combinatorial) theorems into the random setting; see, e.g., the survey of Conlon [8]. The next result provides a random analogue of Theorem 1. Note that it is implicit in the literature (e.g., it is a simple corollary of Theorem 5.2 from [6] and a removal lemma of Green [11, Theorem 1.5]; see also [9, 20, 21]).

Given a finite abelian group G_n , let $G_{n,p}$ be a random subset of G_n obtained by including each element of G_n independently with probability p . Note that throughout the paper $\mathbb{N}$ denotes the set of positive integers (i.e., it does not contain 0).

Theorem 2. *Let $(G_n)_{n \in \mathbb{N}}$ be a sequence of finite abelian groups where $|G_i| < |G_j|$ and $\mu(G_i)/|G_i| = \mu(G_j)/|G_j|$ for all $i < j$. For any $\varepsilon > 0$ there exists $C > 0$ such that the following holds. If $p_n \geq C|G_n|^{-1/2}$ for each $n \in \mathbb{N}$, then*

$$\lim_{n \rightarrow \infty} \mathbb{P}(\text{the largest sum-free set in } G_{n,p_n} \text{ has size } (1 \pm \varepsilon)\mu(G_n) \cdot p_n) = 1.$$

As an illustrative example, a natural class to take in Theorem 2 is $G_n := \mathbb{Z}_q^n = \mathbb{Z}_q \oplus \cdots \oplus \mathbb{Z}_q$ for some fixed prime q . In particular, Theorem 1 implies that $\mu(G_i)/|G_i| = \mu(G_j)/|G_j|$ for all $i, j \in \mathbb{N}$.

Let M be a largest sum-free subset of a finite abelian group G_n . Note that $\mathbb{E}(|G_{n,p} \cap M|) = \mu(G_n) \cdot p$. Thus, Theorem 2 can be interpreted as follows: in a “typical” subset S of G_n of size significantly more than $\sqrt{|G_n|}$, the largest sum-free subset of S has size close to what is “expected”, i.e., close to $|S \cap M|$. Furthermore, it is not difficult to show that the bound on the probability p_n in Theorem 2 is essentially best possible.

Note that there are also related results on the structure of maximum-size sum-free subsets of a random subset of an abelian group G ; see [5, 7].

2. Maximal Sum-Free Subsets

There has been interest in the number of sum-free subsets $f(G)$ of a finite abelian group G . By considering all possible subsets of a largest sum-free subset in G , one obtains that $f(G) \geq 2^{\mu(G)}$. It turns out this trivial bound is not far from

being tight. Indeed, Green and Ruzsa [12] proved that $f(G) = 2^{\mu(G)+o(n)}$ for all abelian groups G of order n . A refined version of this theorem for Type I groups was obtained by Alon, Balogh, Morris and Samotij [1]. Specifically, for such Type I groups they asymptotically determine the number of sum-free sets of a given size m , for all not too small m .

We now consider the *maximal* sum-free version of this problem. That is, we consider the number of maximal sum-free subsets $f_{\max}(G)$ in a finite abelian group G . This problem is wide open in general, though there now are several results on this topic. Improving an earlier bound of Wolfowitz [23], in [4] it was proven that $f_{\max}(G) \leq 3^{\mu(G)/3+o(n)}$ for all abelian groups G of order n ; in particular, this shows that $f_{\max}(G) \ll f(G)$ for all sufficiently large abelian groups. Moreover, Liu and Sharifzadeh [16] showed that $f_{\max}(G) = 3^{\mu(G)/3+o(n)}$ for all Type II groups G whose order n is divisible by 9. In contrast, there are groups where this upper bound is far from tight. In [4] it was shown that $f_{\max}(\mathbb{Z}_2^k) = 2^{\mu(G)/2+o(n)}$, and this bound was further refined in [13] (see Theorem 3 below). Furthermore, in [16] it was shown that there is a $c > 0$ such that almost all even order abelian groups G satisfy $f_{\max}(G) \leq 2^{(1/2-c)\mu(G)}$. Constructions from [4, Proposition 5.7] and [13, Proposition 5.3] show that $f_{\max}(G) \geq 2^{\mu(G)/2-2}$ for all Type III abelian groups whose exponent is 7, 13 or 19. See [4, 13, 16] for further results on $f_{\max}(G)$.

These results paint a rather mixed picture, and it remains unclear how $f_{\max}(G)$ behaves in general. However, the work in [4, 16] implicitly raises the following question.

Question 1. Given an abelian group G of order n , is it true that either $f_{\max}(G) = 3^{\mu(G)/3+o(n)}$ or $f_{\max}(G) \leq 2^{\mu(G)/2+o(n)}$?

Remark 1. Since we submitted this paper, Question 1 has been answered in the negative by Balogh, Garcia, Liu, and Yang [2]. In fact, they have tackled a couple of the other open problems stated in this paper; we further discuss this at the relevant points.

For proving upper bounds on $f_{\max}(G)$ there is now a somewhat well-trodden approach that makes use of the *container method*; see [3, 4, 13, 16]. We will use this approach later in the paper also. Using this approach, in [13] we proved the following sharp result.

Theorem 3 ([13]). *If $k \in \mathbb{N}$ and $n := 2^k$, then $f_{\max}(\mathbb{Z}_2^k) = \left(\binom{n-1}{2} + o(1)\right) 2^{n/4}$. Further, if $k \in \mathbb{N}$ and $n := 3^k$, then $f_{\max}(\mathbb{Z}_3^k) = \left(\frac{(n-3)(n-1)}{3} + o(1)\right) 3^{n/9}$.*

Perhaps the next natural case to consider is $f_{\max}(\mathbb{Z}_5^k)$. On the positive side, a recent result of Lev [15] gives a structural characterisation of the large sum-free subsets of $\mathbb{Z}_5^k$; this result should very likely be needed in order to obtain asymptotically exact bounds on $f_{\max}(\mathbb{Z}_5^k)$. However, despite having this result at hand,

and receiving very nice suggestions from Vsevolod Lev and Wojciech Samotij, we were unable to obtain such sharp bounds. We therefore state the following as a conjecture.

Conjecture 1. If $k \in \mathbb{N}$ and $n := 5^k$, then $f_{\max}(\mathbb{Z}_5^k) = 3^{n/10+o(n)}$.

The following construction shows that the bound on $f_{\max}(\mathbb{Z}_5^k)$ in Conjecture 1 cannot be lowered. Let $B_2 := \{2\} \oplus \mathbb{Z}_5^{k-1}$, and $s := \{1\} \oplus \{0\}$. The map $\phi : b \mapsto -b - s$ is an involution of B_2 , and its only fixed point is $2s$. In particular, there are $(|B_2| - 1)/2$ orbits $\{b, \phi(b)\}$ in B_2 of cardinality 2. Now consider the sets of the form

$$\{s\} \cup \bigcup_{\{b, \phi(b)\} \in B_2/\phi} A_b,$$

where the union ranges over all orbits of cardinality 2 of ϕ , and for each such orbit $\{b, \phi(b)\}$, we choose A_b to be one of $\{b, -b\}$, $\{b, -b-s\}$ or $\{b+s, -b-s\}$. This choice ensures that one can generate $3^{(|B_2|-1)/2} = 3^{n/10-1/2}$ sets S of this form. Further, each such set S is sum-free; this follows as $S \setminus \{s\} \subseteq \{2, 3\} \oplus \mathbb{Z}_5^{k-1}$ is sum-free and the only Schur triples (other than $s, s, 2s$) in $\{s\} \cup (\{2, 3\} \oplus \mathbb{Z}_5^{k-1})$ contain precisely two of $\{b, -b, b+s, -b-s\}$ for some $b \in B_2$. Moreover, one can check that each of these sets lies in a different maximal sum-free subset of $\mathbb{Z}_5^k$. Indeed, given two distinct such sets S, S' the choice of A_b for S and S' must differ for at least one $\{b, \phi(b)\} \in B_2/\phi$. But in this case $S \cup S'$ will contain a Schur triple (formed using s and two of $b, -b, b+s$ and $-b-s$). This proves that $f_{\max}(\mathbb{Z}_5^k) \geq 3^{n/10-1/2}$.

Recently, further structural results for large sum-free subsets of $\mathbb{Z}_p^k$ were obtained when p is a fixed prime where $p \equiv 2 \pmod{3}$; see [19, 22]. These results may also be useful for the $f_{\max}(\mathbb{Z}_p^k)$ problem more generally, although new ideas will again be needed to obtain sharp results.

3. Maximal Distinct Sum-Free Subsets

Let G be an abelian group of order n , and let $f^*(G)$ denote the number of distinct sum-free subsets in G . Let $f_{\max}^*(G)$ denote the number of maximal distinct sum-free subsets in G . As pointed out in [13], the removal lemma of Green [11] implies that $\mu(G) \leq \mu^*(G) \leq \mu(G) + o(n)$ and $f(G) \leq f^*(G) \leq 2^{o(n)}f(G)$. By arguing as in the proof that $f_{\max}(G) \leq 3^{\mu(G)/3+o(n)}$ for all abelian groups G of order n [4, Proposition 5.1], one can also conclude that $f_{\max}^*(G) \leq 3^{\mu(G)/3+o(n)}$. However, we do not know whether $f_{\max}(G)$ is bounded from above by the number of maximal distinct sum-free subsets $f_{\max}^*(G)$ of G for *all* finite abelian groups G . On the other hand, in [13, Section 6.3] we showed that there are abelian groups G for which $f_{\max}^*(G)$ is exponentially larger than $f_{\max}(G)$.

At least for Type I abelian groups, we believe we have a clearer picture as to the behavior of $f_{\max}^*(G)$. Indeed, we conjecture the following.

Conjecture 2. Suppose that G is a Type I abelian group of order n . Then $f_{\max}^*(G) = 2^{\mu(G)/2+o(n)}$.

In [13] we established the $G = \mathbb{Z}_2^k$ case of Conjecture 2. In fact, we showed that $f_{\max}(\mathbb{Z}_2^k) = f_{\max}^*(\mathbb{Z}_2^k)$, and so the result follows by Theorem 3.

The next construction shows that, asymptotically, the bound on $f_{\max}^*(G)$ in Conjecture 2 cannot be lowered in the case of Type I(p) abelian groups for $p \geq 5$. In Section 5, we give constructions that show the bound in Conjecture 2 cannot be lowered in the case of even order groups ($p = 2$); see Proposition 2.

Proposition 1. Suppose $p \geq 5$ and G is a Type I(p) abelian group of order n . Then $f_{\max}^*(G) \geq 2^{\mu(G)/2}$.

Proof. Let B be a sum-free subset of G (so, in particular, a distinct sum-free set) of size $\mu(G) = \frac{p+1}{3p}n$ and such that $B = -B$,¹ e.g.,

$$B := \bigcup_{k=0}^{(p-2)/3} ((3k+1) \cdot g + H),$$

with H a subgroup of G of order $|G|/p$, and $g \notin H$. The map $\sigma : x \mapsto -x$ is an involution of B , and since $p \geq 5$, it has no fixed point. Then, there are $|B|/2$ orbits $\{x, -x\}$ of σ with cardinality 2. Choose one element a_x in each orbit $\{x, -x\}$, and consider the set

$$\{0\} \cup \{a_x \mid \{x, -x\} \in B/\sigma\}.$$

There are $2^{|B|/2} = 2^{\mu(G)/2}$ ways to construct such a set, and notice each such set is distinct sum-free. Further, given two distinct sets S, S' constructed in this way, S and S' must lie in different maximal distinct sum-free subsets of G . Indeed, $S \cup S'$ contains a Schur triple of the form $0, x, -x$. Thus, $f_{\max}^*(G) \geq 2^{\mu(G)/2}$. $\square$

Our next result resolves Conjecture 2 for most even order abelian groups G .

Theorem 4. There exists an absolute constant $C \in \mathbb{N}$ such that the following holds. Let G be an abelian group of even order n , where $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_r}} \oplus K$, with $r, \alpha_1, \dots, \alpha_r \in \mathbb{N}$, and $|K|$ odd. If $n/2^r \geq C$, then

$$f_{\max}^*(G) = 2^{n/4+o(n)}.$$

Remark 2. We now make a number of remarks concerning Theorem 4.

- Note that r may depend on n in the statement of Theorem 4.

¹Actually, Lemma 2 implies that every maximum size sum-free subset B of G satisfies $B = -B$.

- In Section 5, we will see that one can certainly take $C := 10^{10}$.
- The condition that $n/2^r \geq C$ is analogous to a condition in [16, Theorem 3.2], and is a very mild restriction. Indeed, any even order abelian group G not covered by Theorem 4 must be of the form $G = \mathbb{Z}_2^t \oplus H$ for some abelian group H with $|H| < C^2$.
- Theorem 4 tells us that almost all abelian groups G of even order n satisfy $f_{\max}^*(G) = 2^{\mu(G)/2+o(n)}$. This is in contrast to the aforementioned result of Liu and Sharifzadeh [16, Theorem 3.2], which demonstrates that almost all even order abelian groups G satisfy $f_{\max}(G) \ll 2^{\mu(G)/2}$.

In Section 5 we prove – via Theorems 5 and 6 – a sharp version of Theorem 4 that determines the term in front of $2^{n/4}$ rather than having the $o(n)$ term in the exponent.

Whilst Theorem 4 makes substantial progress on the $f_{\max}^*(G)$ problem for even order abelian groups, the picture is still far from clear in general. As a next step, it would be interesting to establish whether the analogue of Question 1 holds for $f_{\max}^*(G)$. In particular, note that by adapting an argument of Liu and Sharifzadeh [16, Proposition 1.5], one can deduce that $f_{\max}^*(G) = 3^{\mu(G)/3+o(n)}$ for all Type II groups G whose order n is divisible by 9.

Remark 3. Since we submitted this paper, Balogh, Garcia, Liu, and Yang [2] have disproved Conjecture 2. Indeed, they have shown that if $G := \mathbb{Z}_2^k \oplus \mathbb{Z}_3$, then $f_{\max}^*(G) = 3^{(1+o(1))\mu(G)/3}$. Further, for every prime $p \geq 23$ with $p \equiv 2 \pmod{3}$, they have shown that there are Type I(p) groups G with $f_{\max}^*(G)$ exponentially larger than $2^{\mu(G)/2}$. On the other hand, they have proven that Conjecture 2 is true for every even order group that is not isomorphic to $\mathbb{Z}_2^k \oplus \mathbb{Z}_3$; see [2, Theorem 1.7]. In light of this, it would be interesting to determine if Conjecture 2 is true for “most” Type I abelian groups.

The rest of the paper is organized as follows. In the next section we introduce various results and concepts that are useful for the proof of Theorem 4, including the notion of a *link graph*. In Section 5 we prove (a strengthening of) Theorem 4 and provide extremal examples for Conjecture 2 for all even order abelian groups. In Section 6 we highlight a problem about maximal independent sets in graphs that we came across when considering Conjecture 1.

4. Useful Results for Theorem 4

4.1. Tools for Abelian Groups

The following is a distinct sum-free subset analogue of a container result of Green and Rusza [12, Proposition 2.1], and is central to our approach.

Lemma 1. *Given any $\delta > 0$ there exists an $n_0 \in \mathbb{N}$ such that the following holds. If G is a finite abelian group of order $n \geq n_0$, then there is a family $\mathcal{F}$ of subsets of G with the following properties.*

- (i) *Every $F \in \mathcal{F}$ has at most δn^2 distinct Schur triples.*
- (ii) *If $S \subseteq G$ is distinct sum-free, then S is contained in some $F \in \mathcal{F}$.*
- (iii) *The cardinality of $\mathcal{F}$ satisfies $|\mathcal{F}| \leq 2^{\delta n}$.*
- (iv) *Every $F \in \mathcal{F}$ is of the form $F = A \cup B$ where A is sum-free and $|B| \leq \delta n$. In particular, $|F| \leq \mu(G) + \delta n$.*

Note that Lemma 1 follows immediately from [11, Theorem 1.5] and, e.g., [6, Theorem 2.2]. We refer to the sets in $\mathcal{F}$ as *containers*. With Lemma 1 at hand, we see that in order to count the number of maximal distinct sum-free sets in an abelian group G , it suffices to count the number of maximal distinct sum-free sets in each container. To do so, the next result provides structure on the sum-free sets in Type I(p) groups.

Lemma 2 ([12]). *Suppose that G is a Type I(p) group and write $p = 3k + 2$. Let $A \subseteq G$ be sum-free, and suppose that $|A| > \left(\frac{1}{3} + \frac{1}{3(p+1)}\right)n$. Then we may find a homomorphism $\psi : G \rightarrow \mathbb{Z}_p$ such that A is contained in $\psi^{-1}(\{k+1, \dots, 2k+1\})$.*

In the proof of Theorem 4, we will use the following easy fact about the number of solutions of the equation $2x = g$ in an even order abelian group.

Fact 1. Let $G := \mathbb{Z}_{2^{\alpha_1}} \oplus \mathbb{Z}_{2^{\alpha_2}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_r}} \oplus K$ where K is an abelian group of odd order and $\alpha_i \in \mathbb{N}$ for each $i \in [r]$. Fix $g \in G$. Then there are at most 2^r solutions in G to the equation $2x = g$.

We end this subsection with a technical lemma that will help us make a sharp count on the number of maximal distinct sum-free subsets in even order abelian groups with few elements of order 2. To prove it, we will use the following well-known correspondence (e.g., it is an immediate corollary of the first proposition in [17, Section 2.2]).

Lemma 3. *Let G be a finite abelian group and $k \in \mathbb{N}$. The number of subgroups of G of order k equals the number of subgroups of G with index k .*

We also require the following fact about subgroups of maximal rank in some even order abelian groups.

Fact 2. Given $\alpha_1 \geq \dots \geq \alpha_r \geq 1$, let $G := \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_r}}$ be a group of rank r . If $H \leq G$ is a subgroup of G , then H has rank r if and only if it contains every order 2 element of G .

Proof. First note that the order 2 elements in G are the non-zero elements of the set $\{0, 2^{\alpha_1-1}\} \oplus \dots \oplus \{0, 2^{\alpha_r-1}\}$, so there are $2^r - 1$ of them. Since $H \leq G$, there exist $\beta_1 \geq \dots \geq \beta_{r'} \geq 1$ such that $H \cong \mathbb{Z}_{2^{\beta_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\beta_{r'}}}$, with $r' \leq r$ the rank of H . Thus, H has $2^{r'} - 1$ order 2 elements. So H contains every order 2 element of G if and only if $2^r - 1 = 2^{r'} - 1$, i.e., if and only if $r = r'$. $\square$

The final lemma of this subsection counts the number of index 2 subgroups of a given rank in some even order abelian groups. Recall that given an abelian group G , $2G$ denotes the subgroup of G whose elements are of the form $2g$ for some $g \in G$.

Lemma 4. *Let r_1 and r_2 be non-negative integers such that $r := r_1 + r_2 \in \mathbb{N}$. Let $G := \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$ with $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$. Then there are $2^{r_1} - 1$ subgroups of index 2 of G with rank r , and $2^r - 2^{r_1}$ subgroups of index 2 of G with rank $r - 1$.*

Proof. Note that a subgroup of index 2 of G has either rank r or $r - 1$. Let $\mathcal{I}_2(G)$ be the set of all the subgroups of G with index 2. Let $\mathcal{R}(G)$ (resp. $\mathcal{R}_{-1}(G)$) be the set of the subgroups of index 2 of G with rank r (resp. $r - 1$). Set $\rho(G) := |\mathcal{R}(G)|$ and $\rho_{-1}(G) := |\mathcal{R}_{-1}(G)|$. By Lemma 3, $|\mathcal{I}_2(G)|$ is equal to the number of subgroups of order 2 of G (which is the number of order 2 elements in G), namely $2^r - 1$. Then, $\rho(G) + \rho_{-1}(G) = 2^r - 1$.

We will now define a bijection φ between $\mathcal{R}(G)$ and $\mathcal{I}_2(2G)$.

Let $H \in \mathcal{R}(G)$, and define $\varphi(H) := 2H$. Let us prove that $\varphi(H) \in \mathcal{I}_2(2G)$. Consider the homomorphism $\pi : G \rightarrow 2G/2H$ defined by $\pi(g) = 2g + 2H$. Then $g \in \ker(\pi)$ if and only if $2g \in 2H$, which is true precisely if (i) $g \in H$ or (ii) there exists $h \in H$ such that $g - h$ has order 2. Note though that in case (ii), $g - h \in H$ since H has rank r and so by Fact 2 it contains every element of G of order 2; this then implies $g \in H$. So in fact $g \in \ker(\pi)$ if and only if $g \in H$ and thus $\ker(\pi) = H$. By the fundamental theorem on homomorphisms, this implies that $2G/2H$ is isomorphic to G/H , proving that $2H \in \mathcal{I}_2(2G)$. This proves that φ is well-defined.

Let $H, H' \in \mathcal{R}(G)$, and suppose that $\varphi(H) = \varphi(H')$, i.e., $2H = 2H'$. Then for every $h \in H$, either $h \in H'$ or there exists $h' \in H'$ such that $h - h'$ has order 2. But since $H' \in \mathcal{R}(G)$, it contains every order 2 element of G by Fact 2; this means $h - h' \in H'$, so $h \in H'$. Thus, $H \subseteq H'$. Similarly $H' \subseteq H$, so $H = H'$, and φ is injective.

Conversely, let $\tilde{H} \in \mathcal{I}_2(2G)$. Then $H := \{g \in G \mid 2g \in \tilde{H}\}$ is such that $\varphi(H) = \tilde{H}$. Indeed, consider the homomorphism $\tilde{\pi} : G \rightarrow 2G/\tilde{H}$ defined by $\tilde{\pi}(g) = 2g + \tilde{H}$. Then by definition $H = \ker(\tilde{\pi})$, so H is an index 2 subgroup of G . Further, H contains every element in G of order 2, and so by Fact 2 must have rank r ; that is, $H \in \mathcal{R}(G)$. This therefore shows φ is surjective.

This bijection φ proves that $\rho(G) = |\mathcal{I}_2(2G)|$. Note that $2G$ has rank r_1 , and so since $2G$ contains $2^{r_1} - 1$ order 2 elements, Lemma 3 implies that $|\mathcal{I}_2(2G)| =$

$2^{r_1} - 1$. Thus, $\rho(G) = 2^{r_1} - 1$, and since $\rho(G) + \rho_{-1}(G) = 2^r - 1$ we also have that $\rho_{-1}(G) = 2^r - 2^{r_1}$. $\square$

4.2. Maximal Independent Sets in Graphs

A key aspect of the approach we take is the translation of the problem of maximal sum-free sets in abelian groups into one about maximal independent sets in graphs. Therefore, we now introduce some notation, and state several results that bound the number of maximal independent sets in graphs.

Let $\Gamma = (V, E)$ be a graph. We define $v(\Gamma) := |V|$ and $e(\Gamma) := |E|$. Given a vertex $x \in V$, we write $d(x, \Gamma)$ for the *degree* of x in Γ (i.e., the number of edges in G incident to x). We write $\delta(\Gamma)$ for the *minimum degree* and $\Delta(\Gamma)$ for the *maximum degree* of Γ . Given a subset $X \subseteq V$ we write $\Gamma \setminus X$ for the subgraph of Γ induced by $V \setminus X$.

We write K_m for the complete graph on m vertices and C_m for the cycle on m vertices. Given graphs Γ and Γ' we write $\Gamma \square \Gamma'$ for the *cartesian product graph*; so its vertex set is $V(\Gamma) \times V(\Gamma')$ and (x, y) and (x', y') are adjacent in $\Gamma \square \Gamma'$ if (i) $x = x'$ and y and y' are adjacent in Γ' or (ii) $y = y'$ and x and x' are adjacent in Γ .

We denote by $\text{MIS}(\Gamma)$ the number of maximal independent sets in Γ . Moon and Moser [18] proved the following bound which holds for any n -vertex graph Γ :

$$\text{MIS}(\Gamma) \leq 3^{n/3}. \quad (1)$$

Note the bound in Equation (1) is tight; consider a graph consisting of the disjoint union of triangles. However, Hujter and Tuza [14] improved this bound in the case of triangle-free n -vertex graphs Γ :

$$\text{MIS}(\Gamma) \leq 2^{n/2}. \quad (2)$$

If Γ is a perfect matching then we have equality in Equation (2). The following lemma improves on Equation (1) in the case of somewhat regular and dense graphs.

Lemma 5 ([3]). *Let $k \geq 1$ and let Γ be a graph on n vertices. Suppose that $\Delta(\Gamma) \leq k\delta(\Gamma)$ and set $b := \sqrt{\delta(\Gamma)}$. Then*

$$\text{MIS}(\Gamma) \leq \sum_{0 \leq i \leq n/b} \binom{n}{i} \cdot 3^{\left(\frac{k}{k+1}\right)\frac{n}{3} + \frac{2n}{3b}}.$$

We will also use the following refined versions of Equations (2) and (1), respectively.

Lemma 6 ([4]). *Let $n, D \in \mathbb{N}$ and $k \in \mathbb{R}$. Suppose that Γ is a graph and T is a set such that $\Gamma' := \Gamma \setminus T$ is triangle-free. Suppose that $\Delta(\Gamma) \leq D$, $v(\Gamma') = n$ and $e(\Gamma') \geq n/2 + k$. Then,*

$$\text{MIS}(\Gamma) \leq 2^{n/2 - k/(100D^2) + 2|T|}.$$

Lemma 7 ([16]). *Let $k \in \mathbb{Z}$, $\Delta \in \mathbb{N}$, and $C := 3^{\Delta/13}$. If Γ is an n -vertex graph with $n + k$ edges and maximum degree Δ , then*

$$\text{MIS}(\Gamma) \leq C \cdot 3^{\frac{n}{3} - \frac{k}{13\Delta}}.$$

The next definition provides the graph that connects our distinct sum-free sets problem to independent sets in graphs.

Definition 2. For subsets $B, S \subseteq G$, let $L_S^*[B]$ be the *distinct link graph of S on B* defined as follows. Its vertex set is B and its edge set consists of the following edges:

- (i) two distinct vertices $x, y \in B$ are adjacent if there exists $s \in S$ such that $\{x, y, s\}$ is a distinct Schur triple;
- (ii) there is a loop at a vertex $x \in B$ if there exist distinct $s, s' \in S$ such that $\{x, s, s'\}$ is a distinct Schur triple.

We call an edge xy in $L_S^*[B]$ a *Type 1* edge if $x - y = s$ for some $s \in S \cup (-S)$, and a *Type 2* edge if it is not a Type 1 edge and $x + y = s$ for some $s \in S$. We denote by $d_i(x, L_S^*[B])$ the number of Type i edges incident to x in $L_S^*[B]$. We write $e_i(L_S^*[B])$ for the number of Type i edges in $L_S^*[B]$.

In Figures 1 and 2 we give a couple of examples of distinct link graphs. For Figure 1, let $S = \{s\}$ such that s has order 3. If $x \in B$ is such that $2x \notin \{0, s, 2s\}$, then s creates 9 edges in the component containing x in $L_S^*[B]$, as shown below. Type 1 edges are represented in blue, and Type 2 edges in red. In Figure 2, note that the set B considered is distinct sum-free but not sum-free.




Figure 1: A component of a distinct link graph.

The following lemma provides the connection between maximal independent sets in the distinct link graph and maximal distinct sum-free sets.

Lemma 8. *Suppose that $B, S \subseteq G$ are both distinct sum-free. If $I \subseteq B$ is such that $S \cup I$ is a maximal distinct sum-free subset of G , then I is a maximal independent set in $L_S^*[B]$.*

Note that the proof of Lemma 8 is identical to the proof [3, Lemma 3.1] (which deals with maximal sum-free subsets of $[n]$).




Figure 2: The distinct link graph $L_{\{s,s'\}}^*[B]$, where s, s' have order 3 and $B = \{s + s', 2s', 2s + 2s'\}$.

5. Proof of Theorem 4

In this section we prove Theorem 4. The proof is based on the general approach initiated in [3, 4], and also the proof adapts parts of arguments that appear in [13, 16]. Note that Theorem 4 follows immediately from the following two results.

Theorem 5 (Sharp upper bound for even order abelian groups). *There exists an absolute constant $C \in \mathbb{N}$ such that the following holds. Given any $\eta > 0$, there exists an $n_0 \in \mathbb{N}$ such that the following holds for all even $n \geq n_0$. Let G be an abelian group of order n , where $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2} \oplus K$, with r_1 and r_2 non-negative integers, $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$, and $|K|$ odd. Set $r := r_1 + r_2 \in \mathbb{N}$. If $n/2^r \geq C$, then*

$$f_{\max}^*(G) \leq (2^{2r-1} + 2^{r+r_1-1} - 2^{r+1} + 2^{r_1})2^{n/4} + (2^r - 2^{r_1} + \eta)2^{n/4-2^{r-2}}.$$

The following theorem shows that the upper bound on $f_{\max}^*(G)$ in Theorem 5 is essentially best possible.

Theorem 6 (Sharp lower bound for even order abelian groups). *There exists an absolute constant $C \in \mathbb{N}$ such that the following holds. Given any $\eta > 0$, there exists an $n_0 \in \mathbb{N}$ such that the following holds for all even $n \geq n_0$. Let G be an abelian group of order n , where $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2} \oplus K$, with r_1 and r_2 non-negative integers, $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$, and $|K|$ odd. Set $r := r_1 + r_2 \in \mathbb{N}$. If $n/2^r \geq C$, then*

$$(2^{2r-1} + 2^{r+r_1-1} - 2^{r+1} + 2^{r_1})2^{n/4} + (2^r - 2^{r_1} - \eta)2^{n/4-2^{r-2}} \leq f_{\max}^*(G).$$

Together Theorems 5 and 6 provide a sharp version of Theorem 4. Though the case $G = \mathbb{Z}_2^k$ is not covered by these two theorems, notice that the coefficient in front of $2^{n/4}$ in Theorems 5 and 6 agrees with coefficient in front of $2^{n/4}$ in Theorem 3 (recall that $f_{\max}^*(\mathbb{Z}_2^k) = f_{\max}(\mathbb{Z}_2^k)$).

The next result provides an extremal example for Conjecture 2 for each even order abelian group.

Proposition 2 (General lower bound for even order abelian groups). *Suppose G is an abelian group of even order n . Then $f_{\max}^*(G) \geq 2^{(n-2)/4}$.*

Proof. Let G be an abelian group of even order n . Thus, $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2} \oplus K$ for some non-negative integers r_1 and r_2 , such that $r := r_1 + r_2 \in \mathbb{N}$, $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$, and $|K|$ is odd. Let H be an index 2 subgroup of $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$ and let $\tilde{A}$ be the coset associated to H that does not contain the identity (i.e., the coset such that $\tilde{A} \neq H$).

We split our argument into two cases.

Case 1: First, suppose that $r_1 \geq 1$ or $r_2 \geq 2$. In this case $|H|$ is even, so there is some $s \in G \setminus (\tilde{A} \oplus K) = H \oplus K$ of order 2. Set $\Gamma := L_{\{0,s\}}^*[\tilde{A} \oplus K]$. We now have two subcases depending on the rank of H .

Subcase 1(a): H has rank $r - 1$. In this subcase, if $x \in \tilde{A} \oplus K$ is such that $2x \notin \{0, s\}$, then the component of Γ containing x is precisely a K_4 containing the vertices $x, x+s, s-x, -x$. If $2x = 0$, then x 's component in Γ is the edge $\{x, x+s\}$; if $2x = s$, then x 's component in Γ is the edge $\{x, -x\}$. Since H has rank $r - 1$, there are 2^{r-1} order 2 elements in $\tilde{A} \oplus K$. Let us denote by $a(s)$ the number of elements $x \in \tilde{A} \oplus K$ such that $2x = s$. Since $\text{MIS}(K_4) = 4$ and $\text{MIS}(K_2) = 2$, we then have

$$\text{MIS}(\Gamma) = 2^{\frac{2^{r-1}+a(s)}{2}} \cdot 4^{\frac{n/2-2^{r-1}-a(s)}{4}} = 2^{n/4}.$$

Subcase 1(b): H has rank r . In this subcase there are no order 2 elements in $\tilde{A} \oplus K$, so Γ is the disjoint union of $a(s)/2$ copies of K_2 and $(n/2 - a(s))/4$ copies of K_4 . Thus,

$$\text{MIS}(\Gamma) = 2^{\frac{a(s)}{2}} \cdot 4^{\frac{n/2-a(s)}{4}} = 2^{n/4}.$$

In both subcases, $\text{MIS}(\Gamma) = 2^{n/4}$. Now if I is a maximal independent set in Γ , then $\{0, s\} \cup I$ is a distinct sum-free subset of G , however, it might not be maximal. But if $I' \neq I$ is another maximal independent set in Γ , then $\{0, s\} \cup I$ and $\{0, s\} \cup I'$ must lie in different maximal distinct sum-free subsets of G ; indeed, otherwise $I \cup I'$ is an independent set in Γ , a contradiction as I and I' are maximal independent sets in Γ . Hence, $f_{\max}^*(G) \geq \text{MIS}(\Gamma) = 2^{n/4}$ in this case.

Case 2: Finally, suppose that $r_1 = 0$ and $r_2 = 1$. Thus, $G = \mathbb{Z}_2 \oplus K$. Set $\Gamma := L_{\{0\}}^*[\{1\} \oplus K]$. In this case, Γ consists of an isolated vertex $\{1\} \oplus \{0\}$, together with a matching where edges are of the form $\{x, -x\}$ for $x \in \{1\} \oplus K$. Thus, $\text{MIS}(\Gamma) = 2^{\frac{|K|-1}{2}} = 2^{(n-2)/4}$. Arguing as in the first case gives $f_{\max}^*(G) \geq \text{MIS}(\Gamma) = 2^{(n-2)/4}$, as desired. $\square$

Note that one can rephrase the proof of Proposition 1 in terms of link graphs as in the proof of Proposition 2.

5.1. Proof of Theorem 5

We now provide a proof of Theorem 5.

Proof of Theorem 5. Let $C := 3235084117$; in footnote 2 we point out the place in the argument where this precise choice of C is used. Given any $\eta > 0$, define additional constants $\delta, \varepsilon, n_0 > 0$ so that

$$0 < 1/n_0 \ll \delta \ll \varepsilon \ll \eta, 1/C.$$

Here the additional constants in the hierarchy are chosen from right to left.

Let G be an abelian group of even order $n \geq n_0$ as in the statement of the theorem. Thus, $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2} \oplus K$ with r_1 and r_2 non-negative integers, $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$, and $|K|$ odd. Furthermore, $n/2^{r_1+r_2} \geq C$.

Apply Lemma 1 to G with parameter δ to obtain a collection $\mathcal{F}$ of containers satisfying conditions (i)–(iv) of the lemma.

By Lemma 1(ii), to obtain an upper bound on $f_{\max}^*(G)$ it suffices to bound the number of maximal distinct sum-free subsets of G that lie in the containers. Given any $F \in \mathcal{F}$, let $f_{\max}^*(F)$ denote the number of maximal distinct sum-free subsets of G that lie in F . Lemma 1(iv) implies that $F = A \cup B$ where A and B are disjoint, A is sum-free and $|B| \leq \delta n$. Thus, each maximal distinct sum-free subset of G lying in F can be obtained in the following way:

- (1) Choose a (perhaps empty) distinct sum-free set S in B ;
- (2) Extend S in A to a maximal one.

This two-step approach, combined with Lemma 8, shows that

$$f_{\max}^*(F) \leq 2^{\delta n} \max_{\substack{S \subseteq B \\ S \text{ is distinct sum-free}}} \text{MIS}(L_S^*[A]),$$

where $L_S^*[A]$ is the distinct link graph of S on A . With this in mind, we now define 7 different types of maximal distinct sum-free subsets of G .

Type 0: Those obtained from a container F where S is chosen to be empty.

Type 1: Those obtained from a container F where $|A| \leq 4n/9$ and $|S| \geq 1$.

Type 2: Those obtained from a container F where $|A| > 4n/9$ and $|S| > 87^2$.

Type 3: Those obtained from a container F where $|A| > 4n/9$ and $2 \leq |S| \leq 87^2$, with $S \neq \{0, s\}$ for every order 2 element $s \in G$.

Type 4: Those obtained from a container F where $|A| > 4n/9$ and $S = \{s\}$ with $s \neq 0$.

Type 5: Those obtained from a container F where $|A| > 4n/9$ and $S = \{0\}$.

Type 6: Those obtained from a container F where $|A| > 4n/9$ and $S = \{0, s\}$ with s an order 2 element.

Let $f_{\max}^{*,i}(G)$ be the total number of maximal distinct sum-free sets of Type i in G . Similarly, we write $f_{\max}^{*,i}(F)$ for the total number of maximal distinct sum-free

sets that lie in the container F . Note that each container F can produce at most one Type 0 maximal distinct sum-free set (namely A), so $f_{\max}^{\star,0}(G) \leq |\mathcal{F}| \leq 2^{\delta n}$ by Lemma 1.

Consider any container $F \in \mathcal{F}$. Now we fix a non-empty distinct sum-free set $S \subseteq B$. In what follows we count how many ways we can extend S in A to a maximal distinct sum-free subset of G . Let $\Gamma := L_S^*[A]$ be the distinct link graph of S on A . From Lemma 8, we see that the number of extensions of S in B to a maximal sum-free set is at most $\text{MIS}(\Gamma)$.

The next claim will be used to bound $f_{\max}^{\star,1}(G)$.

Claim 1. *If $|A| \leq 4n/9$, then $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$.*

To prove Claim 1, by the Moon–Moser bound in Equation (1),

$$\text{MIS}(\Gamma) \leq 3^{|A|/3} \leq 3^{4n/27}.$$

The choice of ε and C implies that $3^{4n/27} \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$ since $n/2^r \geq C$. This completes the proof of Claim 1.

Now we consider Type 2–6 maximal distinct sum-free sets. Suppose $|A| > 4n/9$; by Lemma 2 we can assume that $A \subseteq \tilde{A} \oplus K$ with $\tilde{A}$ a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$. In fact, (by adding elements to the container F if necessary) we may assume $A = \tilde{A} \oplus K$, as $f_{\max}^{\star,i}(F) \leq f_{\max}^{\star,i}(F')$ for $F' \supseteq F$, $i \in [6]$. So A is still sum-free. We can also still assume that A and $B \supseteq S$ are disjoint. Under these conditions, we obtain the following information about the regularity of Γ .

Claim 2. *Let $x \in A$. Then,*

- $d_1(x, \Gamma) = |(S \cup -S) \setminus \{0\}|$;
- $d_2(x, \Gamma) \leq |S|$;
- Γ does not contain any loops.

In particular,

$$|(S \cup -S) \setminus \{0\}| \leq \delta(\Gamma) \leq \Delta(\Gamma) \leq |(S \cup -S) \setminus \{0\}| + |S|.$$

Consequently,

- (i) *if $0 \in S$ and $S = -S$, then $\Delta(\Gamma) \leq 2\delta(\Gamma) + 1$, and*
- (ii) *otherwise, $\Delta(\Gamma) \leq 2\delta(\Gamma)$.*

To prove the first part of Claim 2, note that each element s of $(S \cup -S) \setminus \{0\}$ generates a unique Type 1 neighbor $y = x + s$. For the second part, we have that

x is incident to all edges of the form $(x, s - x)$, with $s \in S$ such that $x \neq s - x$; we have an upper bound because, e.g., some of those edges might be Type 1.

Note that our assumptions ensure $A = g + H$ where H is some index 2 subgroup of G and $g \notin H$, and where $S \subseteq H$. This, together with the definition of the distinct link graph, ensures Γ does not contain loops.

The only possibility for $|(S \cup -S) \setminus \{0\}| + |S|$ to be greater than $2|(S \cup -S) \setminus \{0\}|$ is that $0 \in S$, and $S = -S$. Hence, (i) and (ii) follow. This completes the proof of Claim 2.

Claim 3. *The number of edges in Γ satisfies*

$$e(\Gamma) \geq \frac{|(S \cup -S) \setminus \{0\}| + |S|}{2} |A| - |S|(|S \cup -S| + 1)2^r \geq \frac{\Delta(\Gamma)|A|}{2} - |S|(|S \cup -S| + 1)2^r.$$

By Claim 2, to prove Claim 3 we only need to prove that $e_2(\Gamma) \geq \frac{|S||A|}{2} - |S|(|S \cup -S| + 1)2^r$.

Let X denote the number of pairs $(x, s - x)$ where $x \in A$ and $s \in S$ such that either (i) $(x, s - x)$ is a Type 1 edge in Γ or (ii) $x = s - x$. Note that $e_2(\Gamma) \geq \frac{|S||A|}{2} - X$.

Let $x \in A$ and $s \in S$. Then $(x, s - x)$ is a Type 1 edge in Γ if and only if there exists $s' \in S \cup -S$ such that $x - (s - x) = s'$, or equivalently, $2x = s + s'$. By Fact 1, we deduce there are at most $|S||S \cup -S|2^r$ such edges in Γ .

Similarly, given any $s \in S$, Fact 1 implies that there are at most 2^r solutions $x \in A$ to $x = s - x$. Together with the last paragraph, this implies that $X \leq |S||S \cup -S|2^r + |S|2^r$. This completes the proof of Claim 3.

The next claim will be used to bound $f_{\max}^{\star, 2}(G)$.

Claim 4. *For any $S \subseteq B$ with $|S| > 87^2$, we have $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$.*

To prove Claim 4, by Claim 2 we have $\Delta(\Gamma) \leq 3\delta(\Gamma)$. We apply Lemma 5 with $k = 3$. Noting that $v(\Gamma) = n/2$ and $b := \sqrt{\delta(\Gamma)} \geq 87$, we thus obtain that

$$\text{MIS}(\Gamma) \leq \sum_{0 \leq i \leq n/174} \binom{n/2}{i} 3^{\frac{n}{8} + \frac{n}{261}} \leq \left(\frac{n}{174} + 1\right) (87e)^{\frac{n}{174}} 3^{\frac{n}{8} + \frac{n}{261}}.$$

The choice of C and ε , together with the fact that $n/2^r \geq C$ and n is sufficiently large, ensures that $\left(\frac{n}{174} + 1\right) (87e)^{\frac{n}{174}} 3^{\frac{n}{8} + \frac{n}{261}} \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$, as required. This completes the proof of Claim 4.

The next claim will be used to bound $f_{\max}^{\star, 3}(G)$.

Claim 5. *Consider any distinct sum-free subset $S \subseteq B$ with $2 \leq |S| \leq 87^2$ and $S \neq \{0, s\}$ for every order 2 element $s \in G$. Then $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$.*

To prove Claim 5, our first aim is to show that $\Delta(\Gamma) \geq 4$. We will prove this by considering 8 subcases.

Case 1: $|S| = 4$ and $S = \{0, s_1, s_2, s_3\}$ with s_1, s_2, s_3 having order 2.

Case 2: $|S| = 3$ and $S = \{s_1, s_2, s_3\}$ with s_1, s_2, s_3 having order 2.

Case 3: $|S| = 3$ and $S = \{s_1, s_2, -s_2\}$ with s_1 having order 2.

Case 4: $|S| = 3$ and $S = \{0, s_1, s_2\}$ with s_1 having order 2.

Case 5: $|S| = 2$ and $S = \{s, -s\}$.

Case 6: $|S| = 2$ and $S = \{s_1, s_2\}$ with s_1 having order 2 and $s_2 \neq 0$.

Case 7: $|S| = 2$ and $S = \{0, s\}$ with $s \neq -s$.

Case 8: Any other S satisfying the hypothesis of the claim.

Set $\Delta := \Delta(\Gamma)$. One can check that in **Case 8**, Claim 2 implies that $\Delta \geq |(S \cup -S) \setminus \{0\}| \geq 4$. For example, if $|S| = 4$ and S does not satisfy (i), then $0 \notin S$ or $-S \setminus S \neq \emptyset$ (note that here we are using that S is distinct sum-free). Then Claim 2 ensures that $\Delta \geq 4$.

We can also check by hand that $\Delta \geq 4$ in **Cases 1-7**. Indeed, in **Cases 1 and 2**, consider a vertex $x \in V(\Gamma)$ such that $2x \notin \{s_1, s_1 - s_2, s_1 - s_3, 0\}$; note that such an $x \in V(\Gamma)$ exists by Fact 1 and as $|\Gamma| = n/2 > 4 \cdot 2^r$. Then x has degree at least 4 in Γ since $x + s_1, x + s_2, x + s_3$ and $s_1 - x$ are distinct neighbors of x in Γ . In **Case 3**, consider an $x \in V(\Gamma)$ such that $2x \notin \{s_1, s_1 - s_2, s_1 + s_2, 0\}$; again such an $x \in V(\Gamma)$ exists by Fact 1 and as $|\Gamma| = n/2 > 4 \cdot 2^r$. Then x has degree at least 4 in Γ since $x + s_1, x + s_2, x - s_2$ and $s_1 - x$ are distinct neighbors of x . In **Cases 4 and 6**, consider an $x \in V(\Gamma)$ such that $2x \notin \{0, s_1 - s_2, s_2 - s_1, s_1, s_2\}$; then $x + s_1, x + s_2, s_1 - x$ and $s_2 - x$ are distinct neighbors of x . In **Case 5**, consider an $x \in V(\Gamma)$ such that $2x \notin \{0, -2s, 2s, s, -s\}$; then $x + s, x - s, s - x$ and $-s - x$ are distinct neighbors. Finally, in **Case 7**, consider an $x \in V(\Gamma)$ such that $2x \notin \{2s, 0, s, -s\}$; then $x + s, x - s, -x$ and $s - x$ are distinct neighbors of x .

Now we have proved that $\Delta \geq 4$, we can complete the proof of the claim. By Claim 3 we have $e(\Gamma) \geq \frac{\Delta n}{4} - |S|(|S \cup -S| + 1)2^r$. Applying Lemma 7 with $k := \lceil (\Delta - 2)n/4 - |S|(|S \cup -S| + 1)2^r \rceil$, we have

$$\text{MIS}(\Gamma) \leq 3^{\frac{\Delta}{13}} \cdot 3^{\frac{n}{6} - \frac{(\Delta-2)n}{52\Delta} + \frac{|S|(|S \cup -S| + 1) \cdot 2^r}{13\Delta}}.$$

Since $|S|(|S \cup -S| + 1) \leq 87^2 \cdot (2 \cdot 87^2 + 1)$ and $4 \leq \Delta \leq 3 \cdot 87^2$ by Claim 2, we have

$$\text{MIS}(\Gamma) \leq 3^{\frac{3 \cdot 87^2}{13}} \cdot 3^{\left(\frac{49}{312} \frac{n}{2^r} + \frac{87^2 \cdot (2 \cdot 87^2 + 1)}{52}\right) \cdot 2^r}.$$

The choice of ε and C implies that $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$, since $n/2^r \geq C$.² This completes the proof of Claim 5.

The next claim will be used to bound $f_{\max}^{\star,4}(G)$.

Claim 6. *Let $S = \{s\}$ with $s \neq 0$. Then $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$.*

To prove Claim 6, let ℓ denote the order of s . We first handle the case when $\ell \in \{2, 3\}$, and then $\ell > 3$.

²This is the place in the argument where we use the precise choice of C .

Case (a): $\ell = 2$. In this case, if $x \in V(\Gamma) = A$ is such that $2x \notin \{0, s\}$, then the component of Γ containing x is precisely a 4-cycle $\{x, x+s, -x, s-x\}$. If $x \in V(\Gamma)$ is such that $2x \in \{0, s\}$, the component containing x is an edge $\{x, x+s\}$. Let $I_2 := \{x \in V(\Gamma) \mid 2x \in \{0, s\}\}$. By Fact 1, $|I_2| \leq 2^{r+1}$. Then, $\text{MIS}(\Gamma) \leq 2^{\frac{|A|-|I_2|}{4}} \cdot 2^{|I_2|/2} \leq 2^{n/8+2^{r-1}}$. The choice of ε and C implies that $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$, since $n/2^r \geq C$.

Case (b): $\ell = 3$. In this case, if $x \in V(\Gamma)$ is such that $2x \notin \{0, s, 2s\}$, then the component of Γ containing x is precisely a $C_3 \square K_2$, a perfect matching between the triangles $\{x, x+s, x+2s\}$ and $\{-x+s, -x, -x+2s\}$. If $x \in V(\Gamma)$ such that $2x \in \{0, s, 2s\}$, then the component of Γ containing x is a triangle $\{x, x+s, x+2s\}$. Let $I_3 := \{x \in V(\Gamma) \mid 2x \in \{0, s, 2s\}\}$. By Fact 1, $|I_3| \leq 3 \cdot 2^r$. Noting that $C_3 \square K_2$ contains precisely 6 maximal independent sets, we have that

$$\text{MIS}(\Gamma) \leq 6^{\frac{|A|-|I_3|}{6}} \cdot 3^{|I_3|/3} \leq 6^{n/12-2^{r-1}} \cdot 3^{2^r}.$$

The choice of ε and C implies that $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$, since $n/2^r \geq C$.

Case (c): Now assume $\ell > 3$. Then we can adapt [16, Claim 3.9] to our distinct sum-free set setting: there exists a subset $A_t \subset A = V(\Gamma)$ with $|A_t| \leq 2^r$ that intersects all triangles in Γ . The proof is exactly the same as in the non-distinct setting (see [16, Claim 3.9]), so we omit it. Let A_t be such a set, and let $\Gamma' := \Gamma \setminus A_t$. By Claim 2, every vertex in Γ has degree at most 3, and by Claim 3, $e(\Gamma) \geq \frac{3|A|}{2} - 3 \cdot 2^r$, and thus,

$$e(\Gamma') \geq e(\Gamma) - 3|A_t| \geq \frac{3|A|}{2} - 6 \cdot 2^r.$$

By Lemma 6, with $D := 3$ and $k := |A| - 6 \cdot 2^r$, we obtain

$$\text{MIS}(\Gamma) \leq 2^{\frac{n}{4} - \frac{n/2 - 6 \cdot 2^r}{900} + 2 \cdot 2^r} = 2^{\frac{449}{1800}n + \frac{301}{150} \cdot 2^r}.$$

The choice of ε and C implies that $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$, since $n/2^r \geq C$. This completes the proof of Claim 6.

With the previous claims at hand, it is now straightforward to bound the number of Type 0–4 maximal distinct sum-free subsets of G .

Claim 7. *We have*

$$f_{\max}^{\star,0}(G) + f_{\max}^{\star,1}(G) + f_{\max}^{\star,2}(G) + f_{\max}^{\star,3}(G) + f_{\max}^{\star,4}(G) \leq \eta \cdot 2^{n/4-2^{r-2}}.$$

To prove Claim 7, we already saw that $f_{\max}^{\star,0}(G) \leq 2^{\delta n}$. By Lemma 1, there are at most $2^{\delta n}$ containers $F \in \mathcal{F}$. For each choice of F , there are at most $2^{\delta n}$ choices for $S \subseteq B$. For any S as in the definitions of Types 1–4, Claims 1, 4, 5 and 6 imply that $\text{MIS}(\Gamma) \leq 2^{(1/4-\varepsilon)n-2^{r-2}}$. Thus, by Lemma 8,

$$f_{\max}^{\star,i}(G) \leq 2^{\delta n} \cdot 2^{\delta n} \cdot 2^{(1/4-\varepsilon)n-2^{r-2}} = 2^{(1/4+2\delta-\varepsilon)n-2^{r-2}}$$

for each $i \in [4]$. As $\delta \ll \varepsilon \ll \eta$, Claim 7 now follows.

The next two claims provide an upper bound on the number of Type 5 and 6 maximal distinct sum-free subsets of G .

Claim 8. *We have*

$$f_{\max}^{\star,5}(G) \leq (2^{r_1} - 1)2^{n/4} + (2^r - 2^{r_1})2^{n/4-2^{r-2}}.$$

To prove Claim 8, recall that we assume that each relevant container is of the form $F = A \cup B$ where $A = \tilde{A} \oplus K$ with $\tilde{A}$ a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$. Moreover, for Type 5 maximal distinct sum-free sets, $S = \{0\}$; so to upper bound $f_{\max}^{\star,5}(G)$, it suffices to sum up the number of maximal independent sets in all distinct link graphs of the form $L_{\{0\}}^{\star}[\tilde{A} \oplus K]$. Let $\tilde{A}$ be a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, and set $\Gamma := L_{\{0\}}^{\star}[\tilde{A} \oplus K]$. We distinguish two cases depending on the structure of $\tilde{A}$.

Case (i): $\tilde{A}$ is a coset associated to a subgroup of index 2 with rank $r - 1$. In this case, Γ consists of a set of isolated vertices $I := \{x \in A \mid 2x = 0\}$ together with a matching where edges are of the form $\{x, -x\}$ for $2x \neq 0$. Since $\tilde{A}$ is a coset associated to a subgroup of index 2 with rank $r - 1$, we have $|I| = 2^{r-1}$. Thus, $\text{MIS}(\Gamma) = 2^{\frac{|A| - |I|}{2}} = 2^{n/4-2^{r-2}}$. By Lemma 4, there are $2^r - 2^{r_1}$ such sets $\tilde{A}$; thus such link graphs give rise to at most $(2^r - 2^{r_1})2^{n/4-2^{r-2}}$ Type 5 maximal distinct sum-free sets.

Case (ii): $\tilde{A}$ is a coset associated to a subgroup of index 2 with rank r . In this case, there are no elements in $A = \tilde{A} \oplus K$ such that $2x = 0$. Then, Γ is a perfect matching and $\text{MIS}(\Gamma) = 2^{n/4}$. By Lemma 4, there are $2^{r_1} - 1$ such sets $\tilde{A}$; thus such link graphs give rise to at most $(2^{r_1} - 1)2^{n/4}$ Type 5 maximal distinct sum-free sets.

Summing over all the possible sets A , we obtain $f_{\max}^{\star,5}(G) \leq (2^{r_1} - 1)2^{n/4} + (2^r - 2^{r_1})2^{n/4-2^{r-2}}$. This completes the proof of Claim 8.

Claim 9. *We have*

$$f_{\max}^{\star,6}(G) \leq (2^{2r-1} + 2^{r+r_1-1} - 2^{r+1} + 1)2^{n/4}.$$

To prove Claim 9, as in Claim 8, we assume that each relevant container is of the form $F = A \cup B$ where $A = \tilde{A} \oplus K$ with $\tilde{A}$ a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, and where A and B are disjoint. Moreover, for Type 6 maximal distinct sum-free sets, $S = \{0, s\}$ for some order 2 element $s \in B$; so to upper bound $f_{\max}^{\star,6}(G)$, it suffices to sum up the number of maximal independent sets in all distinct link graphs of the form $L_{\{0,s\}}^{\star}[\tilde{A} \oplus K]$ where $s \notin \tilde{A} \oplus K$ has order 2. Let $\tilde{A}$ be a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, let $s \notin \tilde{A} \oplus K$ be an order 2 element, and set $\Gamma := L_{\{0,s\}}^{\star}[\tilde{A} \oplus K]$. We saw in the proof of Proposition 2 that $\text{MIS}(\Gamma) = 2^{n/4}$. Now let us count the pairs $(\tilde{A}, \{0, s\})$.

Case (i): $\tilde{A}$ is a coset associated to a subgroup of index 2 with rank $r - 1$. By Lemma 4, there are $2^r - 2^{r_1}$ such sets $\tilde{A}$. Since there are $2^{r-1} - 1$ order 2 elements of G not belonging to $A = \tilde{A} \oplus K$, there are $2^{r-1} - 1$ choices for s . Thus, such pairs $(\tilde{A}, \{0, s\})$ contribute at most $(2^r - 2^{r_1})(2^{r-1} - 1)2^{n/4}$ Type 6 maximal distinct sum-free sets.

Case (ii): $\tilde{A}$ is a coset associated to a subgroup of index 2 with rank r . By Lemma 4, there are $2^{r_1} - 1$ such sets $\tilde{A}$. Since $A = \tilde{A} \oplus K$ contains no order 2 elements, there are $2^r - 1$ choices for s . Thus, such pairs $(\tilde{A}, \{0, s\})$ contribute at most $(2^{r_1} - 1)(2^r - 1)2^{n/4}$ Type 6 maximal distinct sum-free sets.

Summing over all the possible choices, we obtain

$$\begin{aligned} f_{\max}^{\star,6}(G) &\leq [(2^r - 2^{r_1})(2^{r-1} - 1) + (2^{r_1} - 1)(2^r - 1)] 2^{n/4} \\ &= (2^{2r-1} + 2^{r+r_1-1} - 2^{r+1} + 1)2^{n/4}. \end{aligned}$$

This completes the proof of Claim 9.

Since every maximal distinct sum-free subset of G is of Type 0–6, Claims 7–9 yield the desired upper bound:

$$f_{\max}^{\star}(G) \leq (2^{2r-1} + 2^{r+r_1-1} - 2^{r+1} + 2^{r_1})2^{n/4} + (2^r - 2^{r_1} + \eta)2^{n/4-2^{r-2}}.$$

□

5.2. Proof of Theorem 6

We now provide a proof of Theorem 6.

Proof of Theorem 6. Let $C := 3235084117$. Given any $\eta > 0$, define additional constants $\varepsilon, n_0 > 0$ so that

$$0 < 1/n_0 \ll \varepsilon \ll \eta, 1/C.$$

Let G be an abelian group of even order $n \geq n_0$ as in the statement of the theorem. Thus, $G = \mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2} \oplus K$ with r_1 and r_2 non-negative integers, $\alpha_1 \geq \dots \geq \alpha_{r_1} \geq 2$, and $|K|$ odd. Furthermore, $n/2^{r_1+r_2} \geq C$.

For a set $A := \tilde{A} \oplus K$ with $\tilde{A}$ a coset (not containing the zero element) associated to a subgroup of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, and a set $S \subseteq G \setminus A$, we say that the pair (A, S) *generates* a distinct sum-free set D in G if there exists a maximal independent set I in the link graph $L_S^*[A]$ such that $D = I \cup S$.

Claim 10. *Consider $A = \tilde{A} \oplus K$ with $\tilde{A}$ a coset (not containing the zero element) of a subgroup H of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$.*

(i) *If H has rank $r - 1$, then $(A, \{0\})$ generates at least*

$$2^{n/4-2^{r-2}} - (n/2 - 2^{r-1}) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}}$$

maximal distinct sum-free sets in G .

(ii) If H has rank r , then $(A, \{0\})$ generates at least

$$2^{n/4} - (n/2 - 2^r) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}}$$

maximal distinct sum-free sets in G .

(iii) If $s \in G \setminus (A \cup \{0\})$ has order 2, $(A, \{0, s\})$ generates at least

$$2^{n/4} - (n/2 - 2) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}}$$

maximal distinct sum-free sets in G .

We now prove each of (i)–(iii) in turn.

To prove Statement (i) of Claim 10, we fix such an A . In the proof of Claim 8 we saw that $L_{\{0\}}^*[A]$ has $2^{n/4-2^{r-2}}$ maximal independent sets. Let $I \subseteq A$ be such a maximal independent set and suppose that $I \cup \{0\}$ is not a maximal distinct sum-free set in G ; call such an I *bad*. Then there exists $s \in G \setminus (A \cup \{0\})$ such that $\{0, s\} \cup I$ is distinct sum-free. So I is a maximal independent set in $L_{\{0,s\}}^*[A]$.

Suppose first that s does not have order 2. By **Case 7** in the proof of Claim 5, there are at most $2^{(1/4-\varepsilon)n-2^{r-2}}$ such sets I .

Now suppose that s has order 2. We saw in Claim 8 that $L_{\{0\}}^*[A]$ consists of a matching between $|A| - 2^{r-1}$ elements, and has 2^{r-1} isolated vertices. So $|I| = (|A| - 2^{r-1})/2 + 2^{r-1} = (|A| + 2^{r-1})/2$. On the other hand, in the proof of Proposition 2, we saw that $L_{\{0,s\}}^*[A]$ consists of the disjoint union of $(|A| - 2^{r-1} - a(s))/4$ copies of K_4 and $(2^{r-1} + a(s))/2$ copies of K_2 , where $a(s)$ is the number of $x \in A$ such that $2x = s$. So $|I| = (|A| + 2^{r-1} + a(s))/4$, which is possible only if $a(s) = |A| + 2^{r-1} = n/2 + 2^{r-1}$. Since $n/2 > n/C \geq 2^r$, such an I cannot exist since $a(s) \leq 2^r$ by Fact 1.

Thus, all bad I s come from s that do not have order 2. The bound in the claim now follows as there are $n/2 - 2^{r-1}$ choices for $s \in G \setminus (A \cup \{0\})$ that do not have order 2. This shows that (i) holds.

To prove Statement (ii) of Claim 10, we saw in the proof of Claim 8 that $L_{\{0\}}^*[A]$ has $2^{n/4}$ maximal independent sets I . In this case, again all bad I s come from s that do not have order 2. Indeed, suppose $s \in G \setminus A$ has order 2. In the proof of Claim 8 we saw maximal independent sets in $L_{\{0\}}^*[A]$ have size $|A|/2$. Meanwhile, in the proof of Proposition 2 we saw that maximal independent sets in $L_{\{0,s\}}^*[A]$ have size $|A|/4 + a(s)/4$. These sizes coincide only if $a(s) = n/2 > n/C \geq 2^r$, which is impossible by Fact 1. The bound in the claim now follows by **Case 7** from the proof of Claim 5 and the fact that there are $n/2 - 2^r$ elements in $G \setminus (A \cup \{0\})$ that do not have order 2. This shows that (ii) holds.

To prove Statement (iii) of Claim 10, we saw in the proof of Proposition 2 that $L_{\{0,s\}}^*[A]$ has $2^{n/4}$ maximal independent sets I . If $I \cup \{0, s\}$ is not a maximal sum-free subset of G , then there exists $s' \in G \setminus (A \cup \{0, s\})$ such that I is a maximal

independent set in $L_{\{0,s,s'\}}^*[A]$. By **Case 4** in the proof of Claim 5, there are at most $2^{(1/4-\varepsilon)n-2^{r-2}}$ such sets I . The bound in the claim now follows as there are $n/2 - 2$ choices for $s' \in G \setminus (A \cup \{0, s\})$. This shows that (iii) holds.

This completes the proof of Claim 10.

Now we need to be careful because we may count the same maximal distinct sum-free set in two different pairs $(A, \{0\})$ and $(A', \{0\})$.

Claim 11. *Consider distinct sets $A = \tilde{A} \oplus K$ and $A' = \tilde{A}' \oplus K$, with $\tilde{A}$ and $\tilde{A}'$ cosets of subgroups of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$. Then there is at most one maximal distinct sum-free subset of G generated by both $(A, \{0\})$ and $(A', \{0\})$.*

To prove Claim 11, note that $|A \cap A'| = |\tilde{A} \cap \tilde{A}'| \leq \mu(G)/2$. Let D be a maximal distinct sum-free subset of G and suppose that it is generated by both $(A, \{0\})$ and $(A', \{0\})$. Then $D = \{0\} \cup I$ with $I \subseteq A \cap A'$.

Case 1: If $\tilde{A}$ and $\tilde{A}'$ are both cosets associated to subgroups of index 2 with rank $r - 1$, then $|I| = \mu(G)/2 + 2^{r-2}$. Indeed, in this case, in the proof of Claim 8 we saw that the link graph $L_{\{0\}}^*[A]$ consists of a matching containing $\mu(G) - 2^{r-1}$ vertices, as well as 2^{r-1} isolated vertices. So the size of a maximal independent set is $(\mu(G) - 2^{r-1})/2 + 2^{r-1} = \mu(G)/2 + 2^{r-2}$. In particular, $|A \cap A'| \geq |I| \geq \mu(G)/2 + 2^{r-2}$, which is impossible.

Case 2: If $\tilde{A}$ and $\tilde{A}'$ are both cosets associated to subgroups of index 2 with rank r , then $|I| = \mu(G)/2$. Indeed we saw in the proof of Claim 8 that $L_{\{0\}}^*[A]$ is a perfect matching. This is only possible if $I = A \cap A'$. So there is at most one possible set D generated by both.

Case 3: If we have one of each, then $|I| = \mu(G)/2 = \mu(G)/2 + 2^{r-2}$, which is impossible.

This completes the proof of Claim 11.

Claim 12. *Consider two sets $A = \tilde{A} \oplus K$ and $A' = \tilde{A}' \oplus K$, with $\tilde{A}$ and $\tilde{A}'$ cosets of subgroups of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, and $s \in G \setminus (A' \cup \{0\})$ with order 2. Then there is no maximal distinct sum-free set generated by both $(A, \{0\})$ and $(A', \{0, s\})$.*

To prove Claim 12, let D be a maximal distinct sum-free set and suppose for a contradiction that it is generated by both $(A, \{0\})$ and $(A', \{0, s\})$. Then $D = \{0\} \cup I = \{0, s\} \cup I'$ with $I \subseteq A$ and $I' \subseteq A'$. In particular, $|I| = |I'| + 1$, and $s \in I \subseteq A$; so A contains an order 2 element, and thus it is associated to a subgroup with rank $r - 1$. In the proof of Claim 8 we saw that in this case we have $|I| = \mu(G)/2 + 2^{r-2}$.

Case 1: If $\tilde{A}'$ is a coset associated to a subgroup of index 2 with rank $r - 1$, then by the proof of Proposition 2, $|I'| = (\mu(G) + 2^{r-1} + a(s))/4$. Since $|I| = |I'| + 1$, this implies that $a(s) = n/2 + 2^{r-1} - 4 > 2^r$, which is impossible by Fact 1.

Case 2: If $\tilde{A}'$ is a coset associated to a subgroup of index 2 with rank r , then by the proof of Proposition 2, $|I'| = (\mu(G) + a(s))/4$. It follows that $a(s) = n/2 + 2^r - 4 > 2^r$, which is again impossible.

This completes the proof of Claim 12.

Claim 13. *Consider two sets $A = \tilde{A} \oplus K$ and $A' = \tilde{A}' \oplus K$, with $\tilde{A}$ and $\tilde{A}'$ cosets of subgroups of index 2 in $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$, and $s \in G \setminus (A \cup \{0\})$, $s' \in G \setminus (A' \cup \{0\})$ both with order 2. Then there are at most $3^{n/12}$ maximal distinct sum-free sets generated by both $(A, \{0, s\})$ and $(A', \{0, s'\})$.*

To prove Claim 13, let D be a maximal distinct sum-free set generated by both $(A, \{0, s\})$ and $(A', \{0, s'\})$. Then $D = \{0, s\} \cup I = \{0, s'\} \cup I'$ with $I \subseteq A$ and $I' \subseteq A'$. Let $D' := D \setminus \{0, s, s'\}$. Then by Lemma 8, D' is a maximal independent set in $L_{\{0, s, s'\}}[A \cap A']$. Since $|A \cap A'| \leq n/4$, by Equation (1) there are at most $3^{n/12}$ choices for D' , and thus for D , as desired. This completes the proof of Claim 13.

Lemma 4 implies that there are $2^r - 2^{r_1}$ choices for $(A, \{0\})$ in Claim 10(i) and $2^{r_1} - 1$ choices for $(A, \{0\})$ in Claim 10(ii). The argument in the proof of Claim 9 implies that there are $2^{2^{r-1}} + 2^{r+r_1-1} - 2^{r+1} + 1$ choices for $(A, \{0, s\})$ in Claim 10(iii). By Lemma 4 there are $2^r - 1$ subgroups of index 2 of $\mathbb{Z}_{2^{\alpha_1}} \oplus \dots \oplus \mathbb{Z}_{2^{\alpha_{r_1}}} \oplus \mathbb{Z}_2^{r_2}$. Thus, there are at most $\binom{2^r-1}{2}$ choices for (A, A') in each of Claims 11–13. Further, there are at most $(2^r)^4$ choices for $(A, \{0, s\})$ and $(A', \{0, s'\})$ in Claim 13. Therefore, Claims 10–13 imply that

$$\begin{aligned} f_{\max}^*(G) &\geq (2^{r_1} - 1) \left(2^{n/4} - (n/2 - 2^r) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}} \right) \\ &\quad + (2^r - 2^{r_1}) \left(2^{n/4-2^{r-2}} - (n/2 - 2^{r-1}) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}} \right) \\ &\quad + (2^{2^{r-1}} + 2^{r+r_1-1} - 2^{r+1} + 1) \left(2^{n/4} - (n/2 - 2) \cdot 2^{(1/4-\varepsilon)n-2^{r-2}} \right) \\ &\quad - \binom{2^r-1}{2} - (2^r)^4 \cdot 3^{n/12}. \end{aligned}$$

As n is sufficiently large, we have that

$$f_{\max}^*(G) \geq (2^{2^{r-1}} + 2^{r+r_1-1} - 2^{r+1} + 2^{r_1})2^{n/4} + (2^r - 2^{r_1} - \eta)2^{n/4-2^{r-2}}.$$

□

Remark 4. Note that we did not try to optimize the value of C in the proof of Theorem 6. With more careful analysis of the link graphs, it may be possible to reduce the value of C to a single-digit number. Theorem 5 should also hold with a much smaller choice of C , although with our argument the choice of C is essentially tight.

6. A Conjecture on Maximal Independent Sets in Graphs with Perfect Matchings

We conclude this paper by raising a conjecture on the number of maximal independent sets in graphs that contain a perfect matching.

Conjecture 3. Let Γ be an n -vertex graph that contains a perfect matching. Then Γ contains at most $2^{n/2}$ maximal independent sets.

Note that we came to this conjecture whilst studying auxiliary (link) graphs for Conjecture 2. A resolution of Conjecture 3 may therefore be helpful for attacking Conjecture 2. Two examples show that the bound in Conjecture 3 cannot be lowered. Indeed, if Γ itself is a perfect matching then Γ contains precisely $2^{n/2}$ maximal independent sets. Similarly, suppose Γ is the disjoint union of $n/6$ copies of the following graph H : H consists of two disjoint triangles joined together by a single edge. Then H contains 8 maximal independent sets, and thus Γ itself contains precisely $8^{n/6} = 2^{n/2}$ maximal independent sets.

Remark 5. Since we submitted this paper, Balogh, Garcia, Liu, and Yang [2] have proven Conjecture 3 in a strengthened form; see [2, Theorem 1.9]. In particular, the resolution of Conjecture 3 is a crucial ingredient to their resolution (in the negative) of Question 1, as well as their proof of Conjecture 2 in the case of even order groups $G \not\cong \mathbb{Z}_2^k \oplus \mathbb{Z}_3$.

Acknowledgements. The second author was supported by EPSRC grants EP/V002279/1 and EP/V048287/1. The authors are grateful to Vsevolod Lev and Wojciech Samotij for discussions relating to Conjecture 1. This research was partially carried out whilst the second author was visiting the Université Bourgogne Europe. This author is grateful for the hospitality he received. The first author thanks Rafik Souanef for pointing out Lemma 4. The authors are also grateful to the referee for their careful review.

References

- [1] N. Alon, J. Balogh, R. Morris, and W. Samotij, Counting sum-free sets in abelian groups, *Israel J. Math.* **199** (2014), 309–344.
- [2] J. Balogh, R.I. Garcia, H. Liu, and N. Yang, Infinitely many groups exhibiting intermediate growth in maximal sum-free sets, preprint, [arXiv:2509.19248](#).
- [3] J. Balogh, H. Liu, M. Sharifzadeh, and A. Treglown, The number of maximal sum-free subsets of integers, *Proc. Amer. Math. Soc.* **143** (2015), 4713–4721.
- [4] J. Balogh, H. Liu, M. Sharifzadeh, and A. Treglown, Sharp bound on the number of maximal sum-free subsets of integers, *J. Euro. Math. Soc.* **20** (2018), 1885–1911.

- [5] J. Balogh, R. Morris, and W. Samotij, Random sum-free subsets of abelian groups, *Isr. J. Math.* **199** (2014), 651–685.
- [6] J. Balogh, R. Morris, and W. Samotij, Independent sets in hypergraphs, *J. Amer. Math. Soc.* **28** (2015), 669–709.
- [7] N. Bushaw, M. Collares Neto, R. Morris, and P. Smith, The sharp threshold for maximum-size sum-free subsets in even-order abelian groups, *Combin. Probab. Comput.* **24** (2015), 609–640.
- [8] D. Conlon, Combinatorial theorems relative to a random set, in *Proceedings of the International Congress of Mathematicians 2014, Vol. 4*, 303–328.
- [9] D. Conlon and W.T. Gowers, Combinatorial theorems in sparse random sets, *Ann. Math.* **84** (2016), 367–454.
- [10] P.H. Diananda and H.P. Yap, Maximal sum-free sets of elements of finite groups, *Proc. Japan Acad.* **45** (1969), 1–5.
- [11] B. Green, A Szemerédi-type regularity lemma in abelian groups, with applications, *Geom. Funct. Anal.* **15** (2005), 340–376.
- [12] B. Green and I. Ruzsa, Sum-free sets in abelian groups, *Israel J. Math.* **147** (2005), 157–189.
- [13] N. Hassler and A. Treglown, On maximal sum-free sets in abelian groups, *Electron. J. Combin.* **29** (2022), #P2.32.
- [14] M. Hujter and Z. Tuza, The number of maximal independent sets in triangle-free graphs, *SIAM J. Discrete Math.* **6** (1993), 284–288.
- [15] V.F. Lev, Large sum-free sets in $\mathbb{Z}_5^n$, *J. Combin. Theory Ser. A* **205** (2024), 105865.
- [16] H. Liu and M. Sharifzadeh, Groups with few maximal sum-free sets, *J. Combin. Theory Ser. A* **177** (2021), 105333.
- [17] L. Mallet-Burgues, Arithmétique des groupes abéliens finis, preprint, [arXiv:2305.01987](https://arxiv.org/abs/2305.01987).
- [18] J.W. Moon and L. Moser, On cliques in graphs, *Israel J. Math.* **3** (1965), 23–28.
- [19] C. Reiher and S. Zotova, Large sum-free sets in finite vector spaces I, preprint, [arXiv:2408.11232](https://arxiv.org/abs/2408.11232).
- [20] D. Saxton and A. Thomason, Hypergraph containers, *Invent. Math.* **201** (2015), 925–992.
- [21] M. Schacht, Extremal results for random discrete structures, *Ann. Math.* **184** (2016), 331–363.
- [22] L. Versteegen, The structure of large sum-free sets in $\mathbb{F}_p^n$, *Q. J. Math.* **75** (2024), 1057–1071.
- [23] G. Wolfowitz, Bounds on the number of maximal sum-free sets, *Euro. J. Combin.* **30** (2009), 1718–1723.